



# Gastrin (ABT329) Mouse mAb

<b>Catalog No</b>	BYab-15174
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Predict react with Mouse, Rat
<b>Applications</b>	IHC
<b>Gene Name</b>	GAST GAS
<b>Protein Name</b>	Gastric mucin 6;Gastric Mucin;Gastric mucin-6;MUC 6;MUC-6;Muc6;MUC6 Fragment;MUC6 mucin;MUC6 mucin Fragment;MUC6_HUMAN;Mucin 6;Mucin 6 gastric;Mucin 6 oligomeric mucus/gel forming;Mucin glycoprotein F
<b>Immunogen</b>	Synthesized peptide derived from human Gastrin
<b>Specificity</b>	The antibody can specifically recognize human Gastrin protein.
<b>Formulation</b>	PBS, pH7.2, 0.03% Porcolin 300, containing stabilizing protein
<b>Source</b>	Monoclonal Mouse IgG2b, kappa
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
<b>Dilution</b>	IHC-p 1:200-400,
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Gastric mucin 6;Gastric Mucin;Gastric mucin-6;MUC 6;MUC-6;Muc6;MUC6 Fragment;MUC6 mucin;MUC6 mucin Fragment;MUC6_HUMAN;Mucin 6;Mucin 6 gastric;Mucin 6 oligomeric mucus/gel forming;Mucin glycoprotein Fragment;Mucin-6;Secretory mucin MUC6 Fragment
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasmic
<b>Tissue Specificity</b>	Pyloric region of the stomach
<b>Function</b>	function:Gastrin stimulates the stomach mucosa to produce and secrete hydrochloric acid and the pancreas to secrete its digestive enzymes. It also stimulates smooth muscle contraction and increases blood circulation and water secretion in the stomach and intestine.,online information:Gastrin entry,PTM:Sulfation enhances proteolytic processing, and blocks peptide degradation. Levels of sulfation differ between proteolytically-cleaved gastrins. Thus, gastrin-6 is almost 73% sulfated, whereas the larger gastrins are less than

**Nanjing BYabscience technology Co.,Ltd**



50% sulfated. Sulfation levels are also tissue-specific.,PTM:Two different processing pathways probably exist in antral G-cells. In the dominant pathway progastrin is cleaved at three sites resulting in two major bioactive gastrins, gastrin-34 and gastrin-17. In the putative alternative pathway, progastrin may be processed only at the most C-terminal dibasic site resul

#### Background

Gastrin is a hormone whose main function is to stimulate secretion of hydrochloric acid by the gastric mucosa, which results in gastrin formation inhibition. This hormone also acts as a mitogenic factor for gastrointestinal epithelial cells. Gastrin has two biologically active peptide forms, G34 and G17. [provided by RefSeq, Jul 2008],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Human gastric adenocarcinoma tissue was stained with Anti-Gastrin (ABT329) Antibody

**Nanjing BYabscience technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658