



# INI-1 (ABT-INI1) mouse mAb

<b>Catalog No</b>	BYab-15429
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	IHC, WB
<b>Gene Name</b>	SMARCB1 BAF47 INI1 SNF5L1
<b>Protein Name</b>	INI-1
<b>Immunogen</b>	Synthesized peptide derived from human INI-1
<b>Specificity</b>	This antibody detects endogenous levels of human INI-1. Heat-induced epitope retrieval (HIER) Citrate buffer of pH6.0 was highly recommended as antigen repair method in paraffin section
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Mouse, Monoclonal/IgG2b, Kappa
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
<b>Dilution</b>	IHC-p 1:100-500, WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Nucleus.
<b>Tissue Specificity</b>	Bra
<b>Function</b>	disease:Defects in SMARCB1 are a cause of rhabdoid tumor (RDT) [MIM:609322]; also called malignant rhabdoid tumor (MRT). Tumor suppressor. Inactivated in rhabdoid tumors. Rhabdoid tumors are a highly malignant group of neoplasms that usually occur in early childhood. SMARCB1/INI1 is also frequently inactivated in epithelioid sarcomas.,disease:Defects in SMARCB1 are a cause of schwannomatosis [MIM:162091]; also called congenital cutaneous neurilemmomatosis. Schwannomas are benign tumors of the peripheral nerve sheath that usually occur singly in otherwise normal individuals. Multiple schwannomas in the same individual suggest an underlying tumor-predisposition syndrome. The most common such syndrome is NF2. The hallmark of NF2 is the development of bilateral vestibular-nerve schwannomas; but two-thirds or more of

Nanjing BYabscience technology Co.,Ltd



all NF2-affected individuals develop schwannomas in other locations, and der

### Background

The protein encoded by this gene is part of a complex that relieves repressive chromatin structures, allowing the transcriptional machinery to access its targets more effectively. The encoded nuclear protein may also bind to and enhance the DNA joining activity of HIV-1 integrase. This gene has been found to be a tumor suppressor, and mutations in it have been associated with malignant rhabdoid tumors. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2015],


### matters needing attention

Avoid repeated freezing and thawing!


### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Human cerebrum tissue was stained with Anti-INI-1 (ABT-INI1) Antibody



LN18 whole cell lysates were separated by 12% SDS-PAGE, and the membrane was blotted with anti-INI-1(ABT-INI1) antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: LN18 Predicted band size: 44kDa Observed band size: 44kDa