



# NHE-7 Polyclonal Antibody

<b>Catalog No</b>	BYab-16483
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	SLC9A7
<b>Protein Name</b>	Sodium/hydrogen exchanger 7
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human SLC9A7. AA range:531-580
<b>Specificity</b>	NHE-7 Polyclonal Antibody detects endogenous levels of NHE-7 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	SLC9A7; NHE7; Sodium/hydrogen exchanger 7; Na(+)/H(+) exchanger 7; NHE-7; Solute carrier family 9 member 7
<b>Observed Band</b>	80kD
<b>Cell Pathway</b>	Golgi apparatus, trans-Golgi network membrane ; Multi-pass membrane protein . Recycling endosome membrane ; Multi-pass membrane protein . Cell membrane ; Multi-pass membrane protein .
<b>Tissue Specificity</b>	Ubiquitously expressed.
<b>Function</b>	function:Mediates electroneutral exchange of protons for Na(+) and K(+) across endomembranes. May contribute to Golgi volume and cation homeostasis.,miscellaneous:Is not inhibited by amiloride but by benzamil and quinine.,similarity:Belongs to the monovalent cation:proton antiporter 1 (CPA1) transporter (TC 2.A.36) family.,subunit:Interacts with SCAMP1, SCAMP2 and SCAMP5; may participate in its shuttling from trans-Golgi network to recycling endosomes.,tissue specificity:Ubiquitously expressed.,

Nanjing BYabscience technology Co.,Ltd



**Background**

This gene encodes a sodium and potassium/ proton antiporter that is a member of the solute carrier family 9 protein family. The encoded protein is primarily localized to the trans-Golgi network and is involved in maintaining pH homeostasis in organelles along the secretory and endocytic pathways. This protein may enhance cell growth of certain breast tumors. This gene is part of a gene cluster on chromosome Xp11.23. A pseudogene of this gene is found on chromosome 12. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Mar 2012],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from Jurkat and COLO cells, using SLC9A7 Antibody. The lane on the right is blocked with the synthesized peptide.

# SLC9A7-



---

---

---

---

**Nanjing BYabscience technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658