



Cytokeratin 20 mouse mAb(ABT044)

| | |
|---------------------------|--|
| Catalog No | BYab-17643 |
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | IHC;WB;IF |
| Gene Name | KRT20 |
| Protein Name | Cytokeratin-20 |
| Immunogen | Synthesized peptide derived from human CK20 |
| Specificity | The antibody can specifically recognize human CK20 protein, and shows no cross reaction with CK1, 5, 6, 7, 8, 13, 14, 15, 17, 18. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.136% sodium azide. |
| Source | Mouse, Monoclonal/IgG2b, Kappa |
| Purification | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Dilution | IHC-p 1:100-500, WB 1:200-1000, IF 1:100-500 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | Keratin, type I cytoskeletal 20 (Cytokeratin-20;CK-20;Keratin-20;K20;Protein IT) |
| Observed Band | |
| Cell Pathway | Cytoplasm . |
| Tissue Specificity | Expressed predominantly in the intestinal epithelium. Expressed in luminal cells of colonic mucosa. Also expressed in the Merkel cells of keratinized oral mucosa; specifically at the tips of some rete ridges of the gingival mucosa, in the basal layer of the palatal mucosa and in the taste buds of lingual mucosa. |
| Function | developmental stage:First detected at embryonic week 8 in individual 'converted' simple epithelial cells of the developing intestinal mucosa. In later fetal stages, synthesis extends over most goblet cells and a variable number of villus enterocytes. In the developing gastric and intestinal mucosa, expressed in all enterocytes and goblet cells as well as certain 'low-differentiated' columnar cells, whereas the neuroendocrine and Paneth cells are negative.,function:Plays a significant role in maintaining keratin filament organization in intestinal epithelia. When phosphorylated, plays a role in the secretion of mucin in the small intestine.,miscellaneous:There are two types of cytoskeletal and microfibrillar |

Nanjing BYabs science technology Co.,Ltd



keratin: I (acidic; 40-55 kDa) and II (neutral to basic; 56-70 kDa).,PTM:Hyperphosphorylation at Ser-13 occurs during the early stages of apoptosis but becomes less prominent during t

Background

The protein encoded by this gene is a member of the keratin family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. The type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. This cytokeratin is a major cellular protein of mature enterocytes and goblet cells and is specifically expressed in the gastric and intestinal mucosa. The type I cytokeratin genes are clustered in a region of chromosome 17q12-q21. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images




Human appendix tissue was stained with Anti-Cytokeratin 20 (ABT044) Antibody

Nanjing BYabs science technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998


监督电话: 15950492658



Human colon carcinoma tissue was stained with Anti-Cytokeratin 20 (ABT044) Antibody



Human rectal carcinoma tissue was stained with
Anti-Cytokeratin 20 (ABT044) Antibody



SW480 whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-CK20(ABT044) antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: SW480 Predicted band size: 48kDa Observed band size: 48kDa