



ANKT Polyclonal Antibody

Catalog No	BYab-00298
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	NUSAP1
Protein Name	Nucleolar and spindle-associated protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from human NUSAP1. AA range:392-441
Specificity	ANKT Polyclonal Antibody detects endogenous levels of ANKT protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NUSAP1; ANKT; BM-037; Nucleolar and spindle-associated protein 1; NuSAP
Observed Band	49kD
Cell Pathway	Cytoplasm . Nucleus, nucleolus . Cytoplasm, cytoskeleton, spindle . Chromosome . Found in the cytoplasm and nucleolus during interphase and redistributes to the mitotic spindle in prometaphase (By similarity). Localizes to the mitotic spindle during anaphase and telophase then disappears from around the chromosomes during cytokinesis (By similarity). Localizes to multiple distinct regions of chromosomes throughout mitosis. .
Tissue Specificity	Bone marrow,Cervix,Epithelium,Fetal liver,Liver,Lung,Lymph,
Function	domain:The KEN box is required for the FZR1-dependent degradation of this protein subsequent to ubiquitination.,function:Microtubule-associated protein with the capacity to bundle and stabilize microtubules (By similarity). May associate with chromosomes and promote the organization of mitotic spindle microtubules around them.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Ubiquitinated. Ubiquitination by FZR1 may lead to proteasome-dependent degradation of this protein.,similarity:Belongs to the

Nanjing BYabscience technology Co.,Ltd



NUSAP family.,subcellular location:Found in the cytoplasm and nucleolus during interphase and redistributes to the mitotic spindle in prometaphase (By similarity). Localizes to the mitotic spindle during anaphase and telophase then disappears from around the chromosomes during cytokinesis (By similarity). Localizes to multiple distinct regions of chromosomes throughout mitosis.,

Background

NUSAP1 is a nucleolar-spindle-associated protein that plays a role in spindle microtubule organization (Raemaekers et al., 2003 [PubMed 12963707]).[supplied by OMIM, Jun 2009],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of various cells using ANKT Polyclonal Antibody



Western blot analysis of lysates from HT-29 cells, using NUSAP1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using NUSAP1 antibody.