



Elongin A2/3 Polyclonal Antibody

Catalog No	BYab-00386
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IF;ELISA
Gene Name	TCEB3B/TCEB3C
Protein Name	RNA polymerase II transcription factor SIII subunit A2/3
Immunogen	The antiserum was produced against synthesized peptide derived from human ELOA2. AA range:101-150
Specificity	Elongin A2/3 Polyclonal Antibody detects endogenous levels of Elongin A2/3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TCEB3B; TCEB3L; RNA polymerase II transcription factor SIII subunit A2; Elongin-A2; EloA2; Transcription elongation factor B polypeptide 3B; TCEB3C; TCEB3L2; RNA polymerase II transcription factor SIII subunit A3; Elongin-A3; EloA3; Transcr
Observed Band	
Cell Pathway	Nucleus .
Tissue Specificity	Specifically expressed in testis.
Function	domain:The elongin BC complex binding domain is also known as BC-box with the consensus [APST]-L-x(3)-C-x(3)-[AIVL].,function:SIII, also known as elongin, is a general transcription elongation factor that increases the RNA polymerase II transcription elongation past template-encoded arresting sites. Subunit A2 is transcriptionally active but its transcription activity is not enhanced by binding to the dimeric complex of the SIII regulatory subunits B and C (elongin BC complex).,similarity:Contains 1 TFIIS N-terminal domain.,subunit:Heterotrimer of

Nanjing BYabscience technology Co.,Ltd



an A (A1, A2 or A3), B and C subunit.,tissue specificity:Specifically expressed in testis.,

Background

This gene encodes the transcriptionally active subunit of the SIII (or elongin) transcription elongation factor complex, which also includes two regulatory subunits, elongins B and C. This complex acts to increase the rate of RNA chain elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites along the DNA template. Whereas a related protein with similar function, elongin A, is ubiquitously expressed, the encoded protein is specifically expressed in the testis, suggesting it may have a role in spermatogenesis. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of A549 cells, using ELOA2 Antibody. The picture on the right is blocked with the synthesized peptide.