



PinX1 Polyclonal Antibody

| | |
|---------------------------|---|
| Catalog No | BYab-00494 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | IHC;IF;ELISA |
| Gene Name | PINX1 |
| Protein Name | PIN2/TERF1-interacting telomerase inhibitor 1 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human PINX1. AA range:121-170 |
| Specificity | PinX1 Polyclonal Antibody detects endogenous levels of PinX1 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | PINX1; LPTL; LPTS; PIN2/TERF1-interacting telomerase inhibitor 1; Liver-related putative tumor suppressor; Pin2-interacting protein X1; Protein 67-11-3; TRF1-interacting protein 1 |
| Observed Band | |
| Cell Pathway | Nucleus . Nucleus, nucleolus. Chromosome, telomere. Chromosome, centromere, kinetochore. Localizes in nucleoli, at telomere speckles and to the outer plate of kinetochores. Localization to the kinetochore is mediated by its central region and depends on NDC80 and CENPE. |
| Tissue Specificity | Ubiquitous; expressed at low levels. Not detectable in a number of hepatocarcinoma cell lines. |
| Function | domain:The TID (telomerase inhibiting domain) domain is sufficient to bind TERT and inhibit its activity.,function:Inhibits telomerase activity. May inhibit cell proliferation and act as tumor suppressor.,similarity:Belongs to the PINX1 family.,similarity:Contains 1 G-patch domain.,subcellular location:In nucleoli and at telomere speckles.,subunit:Binds to MCRS1, TERT and TERF1.,tissue specificity:Ubiquitous; expressed at low levels. Not detectable in a number of |

Nanjing BYabscience technology Co.,Ltd



hepatocarcinoma cell lines.,

Background

domain:The TID (telomerase inhibiting domain) domain is sufficient to bind TERT and inhibit its activity.,function:Inhibits telomerase activity. May inhibit cell proliferation and act as tumor suppressor.,similarity:Belongs to the PINX1 family.,similarity:Contains 1 G-patch domain.,subcellular location:In nucleoli and at telomere speckles.,subunit:Binds to MCRS1, TERT and TERF1.,tissue specificity:Ubiquitous; expressed at low levels. Not detectable in a number of hepatocarcinoma cell lines.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).