



Estrogen Receptor α (ABT4R) rabbit mAb

货号	BYab-18056
同位型	IgG
应用	WB; IHC; ELISA
种属	Human;
靶点	ESR1
基因名称	Estrogen receptor (ER) (ER-alpha) (Estradiol receptor) (Nuclear receptor subfamily 3 group A member 1)
蛋白名称	Estrogen receptor (ER) (ER-alpha) (Estradiol receptor) (Nuclear receptor subfamily 3 group A member 1)
免疫原	Synthesized peptide derived from human Estrogen Receptor α . AA range:450-595
特异性	This antibody detects endogenous levels of ESR1
组成	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
来源	Monoclonal, Rabbit IgG1, Kappa
稀释	IHC 1:100-500, WB 1:500-1000, ELISA 1:5000-20000
纯化工艺	Recombinant Expression and Affinity purified
纯化工艺 其他名称	Recombinant Expression and Affinity purified ESR1 ESR NR3A1
纯化工艺 其他名称 分子量	Recombinant Expression and Affinity purified ESR1 ESR NR3A1 66kD
 纯化工艺 其他名称 分子量 背景 	Recombinant Expression and Affinity purified ESR1 ESR NR3A1 66kD estrogen receptor 1(ESR1) Homo sapiens This gene encodes an estrogen receptor, a ligand-activated transcription factor composed of several domains important for hormone binding, DNA binding, and activation of transcription. The protein localizes to the nucleus where it may form a homodimer or a heterodimer with estrogen receptor 2. Estrogen and its receptors are essential for sexual development and reproductive function, but also play a role in other tissues such as bone. Estrogen receptors are also involved in pathological processes including breast cancer, endometrial cancer, and osteoporosis. Alternative promoter usage and alternative splicing result in dozens of transcript variants, but the full-length nature of many of these variants has not been determined. [provided by RefSeq, Mar 2014],

Nanjing BYabscience technology Co.,Ltd

修 博研生物 BYabscience	国内优质抗体供应商 电 2000年 精准的 WB 检测服务 24H 在线服务,欢迎咨询 电子
	between the estrogen receptor (ER) and NF-kappa-B in a cell-type specific manner. Decreases NF-kappa-B DNA-binding activity and inhibits NF-kappa-B-mediated transcription from the IL6 promoter
细胞定位	[Isoform 1]: Nucleus . Cytoplasm . Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . A minor fraction is associated with the inner membrane.; [Isoform 3]: Nucleus. Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Single-pass type I membrane protein. Associated with the inner membrane via palmitoylation (Probable). At least a subset exists as a transmembrane protein with a N-terminal extracellular domain. .; Nucleus. Golgi apparatus. Cell membrane. Colocalizes with ZDHHC7 and ZDHHC21 in the Golgi apparatus where most probably palmitoylation occurs. Associated with the plasma membrane when palmitoylated.
组织表达	Widely expressed (PubMed:10970861). Not expressed in the pituitary gland (PubMed:10970861). ; [Isoform 3]: Widely expressed, however not expressed in the pituitary gland.
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd