



CD8 a (ABT155) mouse mAb

Catalog No	BY-ab-15559
Isotype	IgG
Reactivity	Human
Applications	IHC;ELISA
Gene Name	CD8A MAL
Protein Name	CD8 a
Immunogen	Synthesized peptide derived from human CD8 a AA range: 100-235
Specificity	The antibody can specifically recognize human CD8 protein, including two types of dimer: $\alpha\beta$ heterodimer or $\alpha\alpha$ homodimer.
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Mouse, Monoclonal/IgG2a, kappa
Purification	The antibody was affinity-purified from ascites by affinity-chromatography using specific immunogen.
Dilution	IHC 1:200-400. ELISA 1:500-5000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/ 1 year
Synonyms	T-cell surface glycoprotein CD8 alpha chain (T-lymphocyte differentiation antigen T8/Leu-2;CD antigen CD8a)
Observed Band	
Cell Pathway	Membranous
Tissue Specificity	CD8 on thymus-derived T-cells usually consists of a disulfide-linked alpha/CD8A and a beta/CD8B chain. Less frequently, CD8 can be expressed as a CD8A homodimer. A subset of natural killer cells, memory T-cells, intraepithelial lymphocytes, monocytes and dendritic cells expresses CD8A homodimers. Expressed at the cell surface of plasmacytoid dendritic cells upon herpes simplex virus-1 stimulation.


Nanjing BYabs science technology Co.,Ltd




Function	<p>disease:Defects in CD8A are a cause of familial CD8 deficiency (CD8 deficiency) [MIM:608957]. Familial CD8 deficiency is a novel autosomal recessive immunologic defect characterized by absence of CD8+ cells, leading to recurrent bacterial infections.,function:Identifies cytotoxic/suppressor T-cells that interact with MHC class I bearing targets. CD8 is thought to play a role in the process of T-cell mediated killing. CD8 alpha chains binds to class I MHC molecules alpha-3 domains.,online information:CD8 entry,online information:CD8A mutation db,PTM:All of the five most carboxyl-terminal cysteines form inter-chain disulfide bonds in dimers and higher multimers, while the four N-terminal cysteines do not.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:In general heterodimer of an alpha and a beta chain linked by two disulfide bonds. Can also form homodimers. Sho</p>
Background	<p>The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011],</p>
matters needing attention	<p>Avoid repeated freezing and thawing!</p>
Usage suggestions	<p>This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.</p>

Products Images


Nanjing BYabscience technology Co.,Ltd



Human burkitt lymphoma tissue was stained with Anti-CD8 (ABT155) Antibody




Human lymphoma tissue was stained with Anti-CD8 (ABT155) Antibody.



Human tonsil tissue was stained with Anti-CD8 (ABT155) Antibody

Nanjing BYabscience technology Co.,Ltd



Human lymphoma tissue was stained with Anti-CD8 (ABT155) Antibody. Polymer HRP Goat Anti Mouse/Rabbit IgG(H+L) Antibody was used at 37°C 30 minutes.

Nanjing BYabscience technology Co.,Ltd