



## Smad2/3 (Acetyl Lys420/378) mouse mAb

Catalog No	BYmab-00920
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SMAD2 MADH2 MADR2
Protein Name	Smad2/3 (Acetyl Lys420/378)
Immunogen	Synthesized peptide derived from human Smad2/3 (Acetyl Lys420/378)
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat Smad2/3 (Acetyl Lys420/378)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
	5
Purity	≥90%
Purity Storage Stability	
•	≥90%
Storage Stability	≥90% -20°C/1 year Mothers against decapentaplegic homolog 2 (MAD homolog 2;Mothers against DPP homolog 2;JV18-1;Mad-related protein 2;hMAD-2;SMAD family member
Storage Stability Synonyms	≥90% -20°C/1 year Mothers against decapentaplegic homolog 2 (MAD homolog 2;Mothers against DPP homolog 2;JV18-1;Mad-related protein 2;hMAD-2;SMAD family member 2;SMAD 2;Smad2;hSMAD2)
Storage Stability Synonyms Observed Band	<ul> <li>≥90%</li> <li>-20°C/1 year</li> <li>Mothers against decapentaplegic homolog 2 (MAD homolog 2;Mothers against DPP homolog 2;JV18-1;Mad-related protein 2;hMAD-2;SMAD family member 2;SMAD 2;Smad2;hSMAD2)</li> <li>55kD</li> <li>Cytoplasm . Nucleus . Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4 (PubMed:9865696, PubMed:21145499). On dephosphorylation by phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101, PubMed:19289081). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast</li> </ul>
Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Mothers against decapentaplegic homolog 2 (MAD homolog 2;Mothers against DPP homolog 2;JV18-1;Mad-related protein 2;hMAD-2;SMAD family member 2;SMAD 2;Smad2;hSMAD2) 55kD Cytoplasm . Nucleus . Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4 (PubMed:9865696, PubMed:21145499). On dephosphorylation by phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101, PubMed:19289081). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity)

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务, 欢迎咨询



Background	transcription, DNA-dependent, regulation of transcription from RNA polymerase II promoter, RNA processing, protein complex assembly, protein amino acid phosphorylation, phosphorus metabolic process, phosphate metabolic process,cell surface receptor linked signal transduction, enzyme linked receptor protein signaling pathway, transmembrane receptor protein serine/threonine kinase signaling pathway, transforming growth factor beta receptor signaling pathway, SMAD protein complex assembly, intracellular signaling cascade, cell-cell signaling, zygotic determination of dorsal/ventral axis, gastrulation, pattern specification process, mesoderm development, heart	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
Products Images		

Nanjing BYabscience technology Co.,Ltd