



UBF Monoclonal Antibody

Catalog No	BYmab-02147
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	UBTF
Protein Name	Nucleolar transcription factor 1
Immunogen	Synthesized peptide derived from UBF . at AA range: 420-500
Specificity	UBF Monoclonal Antibody detects endogenous levels of UBF protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	UBTF; UBF; UBF1; Nucleolar transcription factor 1; Autoantigen NOR-90; Upstream-binding factor 1; UBF-1
Observed Band	89kD
Cell Pathway	Nucleus, nucleolus .
Tissue Specificity	Lymph,Placenta,Testis,
Function	function:Recognizes the ribosomal RNA gene promoter and activates transcription mediated by RNA polymerase I through cooperative interactions with the species-specific factor SL1. It binds specifically to the upstream control element.,PTM:Phosphorylated and activated by PIK3CA.,similarity:Contains 6 HMG box DNA-binding domains.,subunit:Homodimer. Binds to IRS1 and PIK3CA.,
Background	This gene encodes a member of the HMG-box DNA-binding protein family. The encoded protein plays a critical role in ribosomal RNA transcription as a key component of the pre-initiation complex, mediating the recruitment of RNA polymerase I to rDNA promoter regions. The encoded protein may also play
	Naniing BYabscience technology CoLtd

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

24H 在线服务, 欢迎咨询



Products Images		
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
matters needing attention	Avoid repeated freezing and thawing!	
	important roles in chromatin remodeling and pre-rRNA processing, and its activity is regulated by both phosphorylation and acetylation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Pseudogenes of this gene are located on the short arm of chromosomes 3, 11 and X and the long arm of chromosome 11. [provided by RefSeq, Aug 2011],	

Nanjing BYabscience technology Co.,Ltd