



## C-Jun (Hydroxylated-p244) mouse mAb

BYmab-02272
IgG
Human;Mouse;Rat
WB
JUN
C-Jun (Hydroxylated-p244); Transcription factor AP-1;jun;c-jun; AP-1
Synthesized peptide derived from human C-Jun (Hydroxylated-p244)
This antibody detects endogenous levels of Human,Mouse,Rat C-Jun (Hydroxylated-p244)
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
Transcription factor AP-1 (Activator protein 1;AP1;Proto-oncogene c-Jun;V-jun avian sarcoma virus 17 oncogene homolog;p39)
39-42kD
Niveleus
Nucleus.
Expressed in the developing and adult prostate and prostate cancer cells.

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	organization, cell surface receptor linked signal transduction, enzyme linked receptor protein signaling pathway
Background	function:Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'.,PTM:Phosphorylation enhances the transcriptional activity. Phosphorylated by PRKDC.,similarity:Belongs to the bZIP family.,similarity:Belongs to the bZIP family. Jun subfamily.,similarity:Contains 1 bZIP domain.,subunit:Heterodimer with either FOS or BATF3. Interacts with HIVEP3 (By similarity). Interacts with SMAD3/SMAD4 heterodimers. Interacts with MYBBP1A, SPIB and TCF20. Interacts with COPS5; indirectly leading to its phosphorylation. Interacts with DSIPI; this interaction inhibits the binding of active AP1 to its target DNA.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658