



Kallikrein-1 (Cleaved-Ile25) mouse mAb

	DV 1 00000
Catalog No	BYmab-02308
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	KLK1
Protein Name	Kallikrein-1 (Cleaved-Ile25)
Immunogen	Synthesized peptide derived from human Kallikrein-1 (Cleaved-Ile25)
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat Kallikrein-1 (Cleaved-Ile25, protein was cleaved amino acid sequence between 24-25)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Kallikrein-1 (EC 3.4.21.35;Kidney/pancreas/salivary gland kallikrein;Tissue kallikrein)
Observed Band	25kD
Cell Pathway	
Tissue Specificity	Isoform 2 is expressed in pancreas, salivary glands, kidney, colon, prostate gland, testis, spleen and the colon adenocarcinoma cell line T84.
Function	catalytic activity:Preferential cleavage of Arg- -Xaa bonds in small molecule substrates. Highly selective action to release kallidin (lysyl-bradykinin) from kininogen involves hydrolysis of Met- -Xaa or Leu- -Xaa.,function:Glandular kallikreins cleave Met-Lys and Arg-Ser bonds in kininogen to release Lys-bradykinin.,online information:Kallikrein entry,PTM:The O-linked polysaccharides on Ser-93, Ser-104 and Ser-167 are probably the mucin type linked to GalNAc. In PubMed:3163150, GalNAc was detected with the corresponding peptides but not located.,similarity:Belongs to the peptidase S1 family.,similarity:Belongs to the peptidase S1 family.,similarity:Contains 1 peptidase S1 domain.,tissue specificity:Isoform 2 is expressed in pancreas, salivary glands, kidney, colon, prostate gland, testis,

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



spleen and the colon adenocarcinoma cell line T84.,

Background

Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. This protein is functionally conserved in its capacity to release the vasoactive peptide, Lys-bradykinin, from low molecular weight kininogen. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd