



## PDK1 (Phospho Tyr243) mouse mAb

Catalog No	BYmab-02441
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PDK1 PDHK1
Protein Name	PDHK1 (Phospho Tyr243)
Immunogen	Synthesized peptide derived from human PDHK1 (Phospho Tyr243)
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat PDHK1 (Phospho Tyr243)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	[Pyruvate dehydrogenase [lipoamide]] kinase isozyme 1, mitochondrial (EC 2.7.11.2;Pyruvate dehydrogenase kinase isoform 1)
Observed Band	50kD
Cell Pathway	Mitochondrion matrix .
Tissue Specificity	Expressed predominantly in the heart. Detected at lower levels in liver, skeletal muscle and pancreas.
Function	catalytic activity:ATP + [pyruvate dehydrogenase (acetyl-transferring)] = ADP + [pyruvate dehydrogenase (acetyl-transferring)] phosphate.,function:Inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism.,similarity:Belongs to the PDK/BCKDK protein kinase family.,similarity:Contains 1 histidine kinase domain.,tissue specificity:Expressed predominantly in the heart.,
Background	Pyruvate dehydrogenase (PDH) is a mitochondrial multienzyme complex that catalyzes the oxidative decarboxylation of pyruvate and is one of the major enzymes responsible for the regulation of homeostasis of carbohydrate fuels in
	Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务,欢迎咨询



	Products Images
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
matters needing attention	Avoid repeated freezing and thawing!
	mammals. The enzymatic activity is regulated by a phosphorylation/dephosphorylation cycle. Phosphorylation of PDH by a specific pyruvate dehydrogenase kinase (PDK) results in inactivation. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2013],

Nanjing BYabscience technology Co.,Ltd