



S2P Monoclonal Antibody

Catalog No	BYmab-02775
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MBTPS2
Protein Name	Membrane-bound transcription factor site-2 protease
Immunogen	The antiserum was produced against synthesized peptide derived from human MBTPS2. AA range:301-350
Specificity	S2P Monoclonal Antibody detects endogenous levels of S2P protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MBTPS2; S2P; Membrane-bound transcription factor site-2 protease; Endopeptidase S2P; Sterol regulatory element-binding proteins intramembrane protease; SREBPs intramembrane protease
Observed Band	
Cell Pathway	Membrane ; Multi-pass membrane protein . Cytoplasm .
Tissue Specificity	Expressed in heart, brain, placenta, lung, liver, muscle, kidney and pancreas.
Function	catalytic activity: Cleaves several transcription factors that are type-2 transmembrane proteins within membrane-spanning domains. Known substrates include sterol regulatory element-binding protein (SREBP) -1, SREBP-2 and forms of the transcriptional activator ATF6. SREBP-2 is cleaved at the site 477-DRSRILL- -CVLTFLCLSFNPLTSLLQWGGA-505. The residues Asn-Pro, 11 residues distal to the site of cleavage in the membrane-spanning domain, are important for cleavage by S2P endopeptidase. Replacement of either of these residues does not prevent cleavage, but there is no cleavage if both of these residues are replaced., cofactor: Binds 1 zinc ion per

Nanjing BYabscience technology Co.,Ltd



subunit.,function:Intramembrane proteolysis of sterol-regulatory element-binding proteins (SREBPs) within the first transmembrane segment thereby releasing the N-terminal segment with a portion of the transmembrane segment attached. Site-2 cleavage com

Background

This gene encodes a intramembrane zinc metalloprotease, which is essential in development. This protease functions in the signal protein activation involved in sterol control of transcription and the ER stress response. Mutations in this gene have been associated with ichthyosis follicularis with atrichia and photophobia (IFAP syndrome); IFAP syndrome has been quantitatively linked to a reduction in cholesterol homeostasis and ER stress response.[provided by RefSeq, Aug 2009],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images