



## CLIP-170 Monoclonal Antibody

Catalog No	BYmab-03102
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CLIP1
Protein Name	CAP-Gly domain-containing linker protein 1
lmmunogen	The antiserum was produced against synthesized peptide derived from human CLIP1. AA range:1291-1340
Specificity	CLIP-170 Monoclonal Antibody detects endogenous levels of CLIP-170 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CLIP1; CYLN1; RSN; CAP-Gly domain-containing linker protein 1; Cytoplasmic linker protein 1; Cytoplasmic linker protein 170 alpha-2; CLIP-170; Reed-Sternberg intermediate filament-associated protein; Restin
Observed Band	161kD
Cell Pathway	Cytoplasm . Cytoplasm, cytoskeleton . Cytoplasmic vesicle membrane ; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle . Localizes to microtubule plus ends (PubMed:21646404, PubMed:17889670). Localizes preferentially to the ends of tyrosinated microtubules (By similarity). Accumulates in plasma membrane regions with ruffling and protrusions. Associates with the membranes of intermediate macropinocytic vesicles (PubMed:12433698).
Tissue Specificity	Detected in dendritic cells (at protein level). Highly expressed in the Reed-Sternberg cells of Hodgkin disease.
Function	function:Seems to be a intermediate filament associated protein that links endocytic vesicles to microtubules.,similarity:Contains 2 CAP-Gly domains.,subcellular location:Associated with the cytoskeleton.,tissue specificity:Highly expressed in the Reed-Sternberg cells of Hodgkin's disease.,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	The protein encoded by this gene links endocytic vesicles to microtubules. This gene is highly expressed in Reed-Sternberg cells of Hodgkin disease. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658