



A Cyclase IV Monoclonal Antibody

Catalog No	BYmab-03672
lsotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ADCY4
Protein Name	Adenylate cyclase type 4
Immunogen	The antiserum was produced against synthesized peptide derived from human ADCY4. AA range:195-244
Specificity	A Cyclase IV Monoclonal Antibody detects endogenous levels of A Cyclase IV protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
eter age etalenty	
Synonyms	ADCY4; Adenylate cyclase type 4; ATP pyrophosphate-lyase 4; Adenylate cyclase type IV; Adenylyl cyclase 4
	ADCY4; Adenylate cyclase type 4; ATP pyrophosphate-lyase 4; Adenylate
Synonyms	ADCY4; Adenylate cyclase type 4; ATP pyrophosphate-lyase 4; Adenylate cyclase type IV; Adenylyl cyclase 4
Synonyms Observed Band	ADCY4; Adenylate cyclase type 4; ATP pyrophosphate-lyase 4; Adenylate cyclase type IV; Adenylyl cyclase 4 120kD
Synonyms Observed Band Cell Pathway	 ADCY4; Adenylate cyclase type 4; ATP pyrophosphate-lyase 4; Adenylate cyclase type IV; Adenylyl cyclase 4 120kD Cell membrane ; Multi-pass membrane protein . Cytoplasm . Detected in the zona glomerulosa and the zona fasciculata in the adrenal gland
Synonyms Observed Band Cell Pathway Tissue Specificity	 ADCY4; Adenylate cyclase type 4; ATP pyrophosphate-lyase 4; Adenylate cyclase type IV; Adenylyl cyclase 4 120kD Cell membrane ; Multi-pass membrane protein . Cytoplasm . Detected in the zona glomerulosa and the zona fasciculata in the adrenal gland (at protein level). catalytic activity:ATP = 3',5'-cyclic AMP + diphosphate.,cofactor:Binds 2 magnesium ions per subunit.,enzyme regulation:Insensitive to calcium/calmodulin. Stimulated by the G protein beta and gamma subunit complex.,function:This is a membrane-bound, calmodulin-insensitive adenylyl cyclase



国内优质抗体供应商


精准的 WB 检测服务

24H 在线服务, 欢迎咨询



	adenylate cyclase 4, along with adenylate cyclases 2 and 3, is expressed in olfactory cilia, suggesting that several different adenylate cyclases may couple to olfactory receptors and that there may be multiple receptor-mediated mechanisms for the generation of cAMP signals. Alternative splicing results in transcript variants. [provided by RefSeq, Nov 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd