



# A Cyclase IX Monoclonal Antibody


<b>Catalog No</b>	BYmab-03673
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	ADCY9
<b>Protein Name</b>	Adenylate cyclase type 9
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human ADCY9. AA range:137-186
<b>Specificity</b>	A Cyclase IX Monoclonal Antibody detects endogenous levels of A Cyclase IX protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	ADCY9; KIAA0520; Adenylate cyclase type 9; ATP pyrophosphate-lyase 9; Adenylate cyclase type IX; Adenylyl cyclase 9
<b>Observed Band</b>	150kD
<b>Cell Pathway</b>	Cell membrane ; Multi-pass membrane protein .
<b>Tissue Specificity</b>	Detected in skeletal muscle, pancreas, lung, heart, kidney, liver, brain and placenta (PubMed:9628827, PubMed:10987815). Expressed in multiple cells of the lung, with expression highest in airway smooth muscle (PubMed:12972952).
<b>Function</b>	catalytic activity:ATP = 3',5'-cyclic AMP + diphosphate.,cofactor:Binds 2 magnesium ions per subunit.,enzyme regulation:Insensitive to calcium/calmodulin, forskolin and somatostatin. Stimulated by beta-adrenergic receptor activation.,function:May play a fundamental role in situations where fine interplay between intracellular calcium and cAMP determines the cellular function. May be a physiologically relevant docking site for calcineurin.,similarity:Belongs to the adenylyl cyclase class-4/guanylyl cyclase family.,similarity:Contains 2 guanylate cyclase domains.,tissue specificity:Expressed in multiple cells of the lung, with expression highest in airway smooth muscle.,

**Nanjing BYabscience technology Co.,Ltd**



<b>Background</b>	Adenylate cyclase is a membrane bound enzyme that catalyses the formation of cyclic AMP from ATP. It is regulated by a family of G protein-coupled receptors, protein kinases, and calcium. The type 9 adenylyl cyclase is a widely distributed adenylyl cyclase, and it is stimulated by beta-adrenergic receptor activation but is insensitive to forskolin, calcium, and somatostatin. [provided by RefSeq, Jul 2008],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using A Cyclase IX Monoclonal Antibody