



# Arginase II Monoclonal Antibody

<b>Catalog No</b>	BYmab-03711
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	ARG2
<b>Protein Name</b>	Arginase-2 mitochondrial
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human ARG2. AA range:305-354
<b>Specificity</b>	Arginase II Monoclonal Antibody detects endogenous levels of Arginase II protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	ARG2; Arginase-2; mitochondrial; Kidney-type arginase; Non-hepatic arginase; Type II arginase
<b>Observed Band</b>	38kD
<b>Cell Pathway</b>	Mitochondrion .
<b>Tissue Specificity</b>	Expressed most strongly in kidney and prostate, much less strongly in the brain, skeletal muscle, placenta, lung, mammary gland, macrophage, uterus, testis and gut, but apparently not in the liver, heart and pancreas. Expressed in activated T cells (PubMed:27745970).
<b>Function</b>	catalytic activity:L-arginine + H(2)O = L-ornithine + urea.,cofactor:Binds 2 manganese ions per subunit.,function:May play a role in the regulation of extra-urea cycle arginine metabolism and also in down-regulation of nitric oxide synthesis. Extrahepatic arginase functions to regulate L-arginine bioavailability to NO synthase. Since NO synthase is found in the penile corpus cavernosum smooth muscle, the clitoral corpus cavernosum and the vagina, arginase II plays a role in both male and female sexual arousal. It is therefore a potential target for the treatment of male and female sexual arousal disorders.,online

**Nanjing BYabscience technology Co.,Ltd**



information: Arginase entry, pathway: Nitrogen metabolism; urea cycle; L-ornithine and urea from L-arginine: step 1/1., similarity: Belongs to the arginase family., subunit: Homotrimer., tissue specificity: Expressed most strongly in kidney and prostate, much less strongly in the brain,

#### Background

Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exists (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. The type II isoform encoded by this gene, is located in the mitochondria and expressed in extra-hepatic tissues, especially kidney. The physiologic role of this isoform is poorly understood; it is thought to play a role in nitric oxide and polyamine metabolism. Transcript variants of the type II gene resulting from the use of alternative polyadenylation sites have been described. [provided by RefSeq, Jul 2008],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using Arginase II Monoclonal Antibody