



## Arrestin-C Monoclonal Antibody

Catalog No         BYmab-03722           Isotype         IgG           Reactivity         Human;Rat;Mouse;           Applications         WB           Gene Name         ARR3           Protein Name         Arrestin-C           Immunogen         The antiserum was produced against synthesized peptide derived from human ARRC. AA range:339-388           Specificity         Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Storage Stability         -20°C/1 year           Observed Band         40kD           Cell Pathway         Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segments.           Tissue Specificity         Inner and outer segments, and the inner plexiform regions of the retina.           Function         Could binds to photoactivated-phosphorylated red/green opsins, similarity; telongs to the arrestin fa		
Reactivity         Human;Rat;Mouse;           Applications         WB           Gene Name         ARR3           Protein Name         Arrestin-C           Immunogen         The antiserum was produced against synthesized peptide derived from human ARRC. AA range:339-388           Specificity         Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin           Observed Band         40kD           Cell Pathway         Photoreceptor inner segment. Cell projection, cilium, photoreceptor outer segment.           Tissue Specificity         Inner and outer segments, and the inner plexiform regions of the retina.           Function         function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated redgreen opsins. similarity:Belongs t	Catalog No	BYmab-03722
Applications  Gene Name  ARR3  Protein Name  Arrestin-C  Immunogen  The antiserum was produced against synthesized peptide derived from human ARRC. AA range:339-388  Specificity  Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band  40kD  Cell Pathway  Photoreceptor inner segment. Cell projection, cilium, photoreceptor outer segment.  Tissue Specificity  Inner and outer segments, and the inner plexiform regions of the retina.  Function  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins, similarity/Belongs to the arrestin family, tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.  Background  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the erreceptor from the heterotrineric G protein, resulting in termination of the G protein-coupled receptors. This binding uncouples the receptor from the heterotrineric G protein also is a part of of the G protein-coupled receptors.	Isotype	IgG
Protein Name Arrestin-C  Immunogen The antiserum was produced against synthesized peptide derived from human ARRC. AA range:339-388  Specificity Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band 40kD  Cell Pathway Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Tissue Specificity Inner and outer segments, and the inner plexiform regions of the retina.  Function function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins, similarity. Belongs to the arrestin family, tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.  Background The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein also is a part of	Reactivity	Human;Rat;Mouse;
Protein Name         Arrestin-C           Immunogen         The antiserum was produced against synthesized peptide derived from human ARRC. AA range:339-388           Specificity         Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin           Observed Band         40kD           Cell Pathway         Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .           Tissue Specificity         Inner and outer segments, and the inner plexiform regions of the retina.           Function         function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated retina-specific signal transduction copies, similarity. Belongs to the arrestin family., tissue specificity: Inner and outer segments, and the inner plexiform regions of the retina.           Background         The pro	Applications	WB
Immunogen  The antiserum was produced against synthesized peptide derived from human ARRC. AA range:339-388  Specificity  Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse,IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band  40kD  Cell Pathway  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment.  Tissue Specificity  Inner and outer segments, and the inner plexiform regions of the retina.  Function  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins, similarity. Belongs to the arrestin family. Lissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.  Background  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in terminotor of the G protein, resulting in terminotor of the G protein resulting in terminotor agonist-activated.	Gene Name	ARR3
ARRC. AA range:339-388  Specificity Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band 40kD  Cell Pathway Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment.  Tissue Specificity Inner and outer segments, and the inner plexiform regions of the retina.  Function Could binds to photoactivated-phosphorylated red/green opsins, similarity. Belongs to the arrestin family., tissue specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins, similarity. Belongs to the arrestin family, tissue specificity: Inner and outer segments, and the inner plexiform regions of the retina.  Background The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor signaling. The encoded protein also is a part of	Protein Name	Arrestin-C
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band 40kD  Cell Pathway Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Tissue Specificity Inner and outer segments, and the inner plexiform regions of the retina.  Function function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins, similarity. Belongs to the arrestin family, tissue specificity. Inner and outer segments, and the inner plexiform regions of the retina.  Background The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Immunogen	
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin           Observed Band         40kD           Cell Pathway         Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .           Tissue Specificity         Inner and outer segments, and the inner plexiform regions of the retina.           Function         function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins. similarity:Belongs to the arrestin family., tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.           Background         The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Specificity	Arrestin-C Monoclonal Antibody detects endogenous levels of Arrestin-C protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band 40kD  Cell Pathway Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Tissue Specificity Inner and outer segments, and the inner plexiform regions of the retina.  Function function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins, similarity:Belongs to the arrestin family, tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.  Background The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band 40kD  Cell Pathway Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Tissue Specificity Inner and outer segments, and the inner plexiform regions of the retina.  Function function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.  Background The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin         Observed Band       40kD         Cell Pathway       Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .         Tissue Specificity       Inner and outer segments, and the inner plexiform regions of the retina.         Function       function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.,         Background       The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band 40kD  Cell Pathway Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Tissue Specificity Inner and outer segments, and the inner plexiform regions of the retina.  Function function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.  Background The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band  40kD  Cell Pathway  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Inner and outer segments, and the inner plexiform regions of the retina.  Function  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.,  Background  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Concentration	1 mg/ml
Synonyms  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  Observed Band  40kD  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Tissue Specificity  Inner and outer segments, and the inner plexiform regions of the retina.  Function  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.,  Background  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of		
arrestin-3; X-arrestin  Observed Band  40kD  Cell Pathway  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Inner and outer segments, and the inner plexiform regions of the retina.  Function  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.,  Background  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Purity	≥90%
Cell Pathway  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Inner and outer segments, and the inner plexiform regions of the retina.  function: May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins., similarity: Belongs to the arrestin family., tissue specificity: Inner and outer segments, and the inner plexiform regions of the retina.,  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of		
Tissue Specificity  Inner and outer segments, and the inner plexiform regions of the retina.  Function  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.,  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Storage Stability	-20°C/1 year  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone
Function  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.,  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Storage Stability Synonyms	-20°C/1 year  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin
Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer segments, and the inner plexiform regions of the retina.,  Background  The protein encoded by this gene is a non-visual arrestin which binds to agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Storage Stability Synonyms Observed Band	-20°C/1 year  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  40kD  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer
agonist-activated, phosphorylated G protein-coupled receptors. This binding uncouples the receptor from the heterotrimeric G protein, resulting in termination of the G protein-coupled receptor signaling. The encoded protein also is a part of	Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  40kD  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .
	Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  ARR3; ARRX; CAR; Arrestin-C; Cone arrestin; C-arrestin; cArr; Retinal cone arrestin-3; X-arrestin  40kD  Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment .  Inner and outer segments, and the inner plexiform regions of the retina.  function:May play a role in an as yet undefined retina-specific signal transduction. Could binds to photoactivated-phosphorylated red/green opsins.,similarity:Belongs to the arrestin family.,tissue specificity:Inner and outer

Nanjing BYabscience technology Co.,Ltd





centrosome function. [provided by RefSeq, May 2016],

matters	needing	
attention		

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



