



Cdc42EP2 Monoclonal Antibody

Catalog No	BYmab-03751
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CDC42EP2
Protein Name	Cdc42 effector protein 2
Immunogen	The antiserum was produced against synthesized peptide derived from human BORG1. AA range:10-59
Specificity	Cdc42EP2 Monoclonal Antibody detects endogenous levels of Cdc42EP2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CDC42EP2; BORG1; CEP2; Cdc42 effector protein 2; Binder of Rho GTPases 1
Observed Band	28kD
Cell Pathway	Endomembrane system ; Peripheral membrane protein . Cytoplasm, cytoskeleton .
Tissue Specificity	Highly expressed in the heart. Weakly expressed in the pancreas and liver.
Function	domain: The CRIB domain mediates interaction with CDC42., function: Probably involved in the organization of the actin cytoskeleton. May act downstream of CDC42 to induce actin filament assembly leading to cell shape changes. Induces pseudopodia formation in fibroblasts in a CDC42-dependent manner., similarity: Belongs to the BORG/CEP family., similarity: Contains 1 CRIB domain., subunit: Interacts with RHOQ and CDC42 in a GTP-dependent manner, and with SEPT7., tissue specificity: Highly expressed in the heart. Weakly expressed in the pancreas and liver.,
Function Background	involved in the organization of the actin cytoskeleton. May act downstream of CDC42 to induce actin filament assembly leading to cell shape changes. Induces pseudopodia formation in fibroblasts in a CDC42-dependent manner.,similarity:Belongs to the BORG/CEP family.,similarity:Contains 1 CRIB domain.,subunit:Interacts with RHOQ and CDC42 in a GTP-dependent manner, and with SEPT7.,tissue specificity:Highly expressed in the heart. Weakly

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



protein encoded by this gene is a member of the Borg family of CDC42 effector proteins. Borg family proteins contain a CRIB (Cdc42/Rac interactive-binding)
proteins. Borg family proteins contain a CRIB (Cdc42/Rac interactive-binding)
domain. They bind to, and negatively regulate the function of CDC42.
Coexpression of this protein with CDC42 suggested a role of this protein in actin
filament assembly and cell shape control. [provided by RefSeq, Aug 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Cdc42EP2 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658