



GAS41 Monoclonal Antibody

Catalog No	BYmab-03890
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	YEATS4
Protein Name	YEATS domain-containing protein 4
Immunogen	The antiserum was produced against synthesized peptide derived from human GAS41. AA range:1-50
Specificity	GAS41 Monoclonal Antibody detects endogenous levels of GAS41 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	YEATS4; GAS41; YEATS domain-containing protein 4; Glioma-amplified sequence 41; Gas41; NuMA-binding protein 1; NuBI-1; NuBI1
Observed Band	40kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in brain, heart, kidney, liver, lung, pancreas, placenta and skeletal muscle.
Function	function:Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage.,similarity:Contains 1 YEATS domain.,subunit:Component of numerous

Nanjing BYabscience technology Co.,Ltd



complexes with chromatin remodeling and histone acetyltransferase activity. Component of the NuA4 histone acetyltransferase complex whi

Background

The protein encoded by this gene is found in the nucleoli. It has high sequence homology to human MLLT1, and yeast and human MLLT3 proteins. Both MLLT1 and MLLT3 proteins belong to a class of transcription factors, indicating that the encoded protein might also represent a transcription factor. This protein is thought to be required for RNA transcription. This gene has been shown to be amplified in tumors. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GAS41 Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd