



## Grap Monoclonal Antibody

Catalog No.	RVmah 02006
Catalog No	BYmab-03906
lsotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GRAP
Protein Name	GRB2-related adapter protein
Immunogen	The antiserum was produced against synthesized peptide derived from human GRAP. AA range:1-50
Specificity	Grap Monoclonal Antibody detects endogenous levels of Grap protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GRAP; GRB2-related adapter protein
Observed Band	28kD
Cell Pathway	Membrane ; Peripheral membrane protein . Cell junction, synapse . Localizes at the presynaptic terminal
Tissue Specificity	Blood,Brain,
Function	function:Couples signals from receptor and cytoplasmic tyrosine kinases to the Ras signaling pathway.,similarity:Belongs to the GRB2/sem-5/DRK family.,similarity:Contains 1 SH2 domain.,similarity:Contains 2 SH3 domains.,subunit:Associates through its SH2 domain with ligand-activated receptors for stem cell factor (KIT) and erythropoietin (EPOR). Also forms a stable complex with the Bcr-Abl oncoprotein. GRAP is associated with the Ras guanine nucleotide exchange factor SOS1, primarily through its N-terminal SH3 domain. Interacts with phosphorylated LAT upon TCR activation. Interacts with SHB.,
Background	This gene encodes a member of the GRB2/Sem5/Drk family and functions as a cytoplasmic signaling protein which contains an SH2 domain flanked by two SH3
	Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商

24H 在线服务,欢迎咨询

精准的 WB 检测服务

	domains. The SH2 domain interacts with ligand-activated receptors for stem cell factor and erythropoietin, and facilitates the formation of a stable complex with the BCR-ABL oncoprotein. This protein also associates with the Ras guanine nucleotide exchange factor SOS1 (son of sevenless homolog 1) through its N-terminal SH3 domain. In general, it couples signals from receptor and cytoplasmic tyrosine kinases to the Ras signaling pathway. [provided by RefSeq, Jul 2012],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using Grap Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd