



## IPP-2 Monoclonal Antibody

Catalog NoBYmab-03931IsotypeIgGReactivityHuman;Mouse;RatApplicationsWBGene NamePPP1R2Protein NameProtein phosphatase inhibitor 2ImmunogenThe antiserum was produced against synthesized peptide derived from hum PPP1R2. AA range:86-135SpecificityIPP-2 Monoclonal Antibody detects endogenous levels of IPP-2 protein.FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgG
Reactivity Human; Mouse; Rat  Applications WB  Gene Name PPP1R2  Protein Name Protein phosphatase inhibitor 2  Immunogen The antiserum was produced against synthesized peptide derived from hum PPP1R2. AA range: 86-135  Specificity IPP-2 Monoclonal Antibody detects endogenous levels of IPP-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Applications  Gene Name  PPP1R2  Protein Name  Protein phosphatase inhibitor 2  Immunogen  The antiserum was produced against synthesized peptide derived from hum PPP1R2. AA range:86-135  Specificity  IPP-2 Monoclonal Antibody detects endogenous levels of IPP-2 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name Protein Name Protein phosphatase inhibitor 2  Immunogen The antiserum was produced against synthesized peptide derived from hum PPP1R2. AA range:86-135  Specificity IPP-2 Monoclonal Antibody detects endogenous levels of IPP-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Protein Name Protein phosphatase inhibitor 2  Immunogen The antiserum was produced against synthesized peptide derived from hum PPP1R2. AA range:86-135  Specificity IPP-2 Monoclonal Antibody detects endogenous levels of IPP-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Immunogen  The antiserum was produced against synthesized peptide derived from hum PPP1R2. AA range:86-135  Specificity  IPP-2 Monoclonal Antibody detects endogenous levels of IPP-2 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
PPP1R2. AA range:86-135  Specificity IPP-2 Monoclonal Antibody detects endogenous levels of IPP-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Managharat Managha
Source Monoclonal, Mouse,IgG
Our of the second of the secon
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b> WB 1:500-2000
Concentration 1 mg/ml
Purity ≥90%
Storage Stability -20°C/1 year
Synonyms PPP1R2; IPP2; Protein phosphatase inhibitor 2; IPP-2
Observed Band 25kD
Cell Pathway protein phosphatase type 1 complex,
Tissue Specificity Epithelium, Heart, Platelet, Skin, Testis,
function: Inhibitor of protein-phosphatase 1.,PTM:Phosphorylation on Thr-73 GSK3 activates PP1 by dissociating the PP1-PPP1R2 complex (By similarity Phosphorylation on Ser-44 by ATM activates PP1 by dissociating the PP1-PPP1R2 complex.,similarity:Belongs to the protein phosphatase inhibit family.,subunit:Heterodimer with PP1.,
Protein phosphatase-1 (PP1) is one of the main eukaryotic serine/threoning phosphatases. The protein encoded by this gene binds to the catalytic subur PP1, strongly inhibiting its activity. Ten related pseudogenes have been four throughout the human genome. Several splice variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015],

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

