



MRP-S34 Monoclonal Antibody

Catalog No BYmab-04018 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name MRPS34 Protein Name 28S ribosomal protein S34 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPS34. AA range:169-218 Specificity MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteims. Background		
Reactivity Applications WB Gene Name MRPS34 Protein Name 28S ribosomal protein S34 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPS34. AA range:169-218 Specificity MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit.Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins. Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes is that the latter contain a 5s rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Catalog No	BYmab-04018
Applications WB Gene Name MRPS34 Protein Name 28S ribosomal protein S34 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPS34. AA range:169-218 Specificity MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins. Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes is that the latter contains and siffer greatly	Isotype	IgG
Gene Name MRPS34 Protein Name 28S ribosomal protein S34 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPS34. AA range:169-218 Specificity MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins. Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrion. Mitochondrial ribosomes in the protein synthesis within the mitochondrion or mitochondrial ribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Reactivity	Human;Mouse
Protein Name 28S ribosomal protein S34 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPS34. AA range:169-218 Specificity MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit: Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrion. Mitochondrial ribosomes consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a SS rRNA. Among different repoteins comprising the mitoribosome differ greatly.	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human MRPS34. AA range:169-218 Specificity MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrion. Intochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosome differ greatly frederien comprising the mitoribosome differ greatly filed proteins some mitoribosome differ greatly filed proteins or proteins suffer greatly filed proteins compared to prokaryotic ribosomes is that the latter contain a SS rRNA. Among different proteins comprising the mitoribosome differ greatly filed proteins compared to prokaryotic ribosomes is that the latter contain a SS rRNA.	Gene Name	MRPS34
MRPS34. AA range:169-218 Specificity MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion. Tissue Specificity Eye, Function subunit: Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins. Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrian Allochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition contain a 5S rRNA. Among different proteins, and in reverse defined the mitoribosomes is reversed. Another difference between mammalian mitoribosomes and proteins for reversed. Another difference between mammalian mitoribosomes and species, the proteins comprising the mitoribosome differ greatly the mitoribosome differ greatly defined the mitoribosome differ greatly defined and the mitoribosome differ greatly defined and protein such protein such a protein such and the mitoribosome differ greatly the proteins comprising the mitoribosome differ greatly defined and protein such and the protein such and the mitoribosome differ greatly defined and protein such and the protein su	Protein Name	28S ribosomal protein S34 mitochondrial
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among difference between mammalian mitoribosome some offere greatly the proteins comprising the mitoribosome differe greatly	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit: Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins. Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Specificity	MRP-S34 Monoclonal Antibody detects endogenous levels of MRP-S34 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes anhelp in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes anhelp in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes an help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion. Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Concentration	1 mg/ml
Synonyms MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt Observed Band 26kD Cell Pathway Mitochondrion . Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosome and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Purity	≥90%
Observed Band Cell Pathway Mitochondrion . Tissue Specificity Eye, Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Storage Stability	-20°C/1 year
Cell Pathway Mitochondrion . Eye, Function Subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Synonyms	MRPS34; 28S ribosomal protein S34; mitochondrial; MRP-S34; S34mt
Tissue Specificity Eye, Subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Observed Band	26kD
Function subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Cell Pathway	Mitochondrion .
Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Tissue Specificity	Eye,
help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly	Function	subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins.,
	Background	(mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly


Nanjing BYabscience technology Co.,Ltd





	recognition by sequence homology. This gene encodes a 28S subunit protein. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Nanjing BYabscience technology Co.,Ltd