



## MRP-S7 Monoclonal Antibody

Catalog No	BYmab-04022
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MRPS7
Protein Name	28S ribosomal protein S7 mitochondrial
Immunogen	The antiserum was produced against synthesized peptide derived from human MRPS7. AA range:91-140
Specificity	MRP-S7 Monoclonal Antibody detects endogenous levels of MRP-S7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MRPS7; 28S ribosomal protein S7; mitochondrial; MRP-S7; S7mt; bMRP-27a; bMRP27a
Observed Band	24kD
Cell Pathway	Mitochondrion .
Tissue Specificity	Eye,Skeletal muscle,Small intestine,Umbilical cord blood,
Function	similarity:Belongs to the ribosomal protein S7P family.,subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins.,
Background	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein. In the prokaryotic ribosome, the comparable protein is thought to play an essential role in organizing the 3' domain of the 16 S rRNA in the vicinity of the P-and A-sites. Pseudogenes corresponding to
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658