



NACAD Monoclonal Antibody

Catalog No	BYmab-04028
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	NACAD
Protein Name	NAC-alpha domain-containing protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from human NACAD. AA range:376-425
Specificity	NACAD Monoclonal Antibody detects endogenous levels of NACAD protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NACAD; KIAA0363; NAC-alpha domain-containing protein 1
Observed Band	140kD
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Brain,
Function	function:May prevent inappropriate targeting of non-secretory polypeptides to the endoplasmic reticulum (ER). May bind to nascent polypeptide chains as they emerge from the ribosome and block their interaction with the signal recognition particle (SRP), which normally targets nascent secretory peptides to the ER. May also reduce the inherent affinity of ribosomes for protein translocation sites in the ER membrane (M sites).,similarity:Belongs to the NAC-alpha family.,similarity:Contains 1 NAC-A/B (NAC-alpha/beta) domain.,
Background	function:May prevent inappropriate targeting of non-secretory polypeptides to the endoplasmic reticulum (ER). May bind to nascent polypeptide chains as they emerge from the ribosome and block their interaction with the signal recognition

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



particle (SRP), which normally tal also reduce the inherent affinity of ER membrane (M sites).,similarity family.,similarity:Contains 1 NAC	rgets nascent secretory peptides to the ER. May if ribosomes for protein translocation sites in the y:Belongs to the NAC-alpha -A/B (NAC-alpha/beta) domain.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NACAD Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658