



# RGS1 Monoclonal Antibody

<b>Catalog No</b>	BYmab-04119
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	RGS1
<b>Protein Name</b>	Regulator of G-protein signaling 1
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human RGS1. AA range:118-167
<b>Specificity</b>	RGS1 Monoclonal Antibody detects endogenous levels of RGS1 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	RGS1; 1R20; BL34; IER1; Regulator of G-protein signaling 1; RGS1; B-cell activation protein BL34; Early response protein 1R20
<b>Observed Band</b>	22kD
<b>Cell Pathway</b>	Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm, cytosol .
<b>Tissue Specificity</b>	Detected in peripheral blood monocytes (PubMed:10480894). Expression is relatively low in B-cells and chronic lymphocytic leukemia B-cells; however, in other types of malignant B-cell such as non-Hodgkin lymphoma and hairy cell leukemia, expression is constitutively high (PubMed:8473738).
<b>Function</b>	function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. This protein may be involved in the regulation of B-cell activation and proliferation.,induction:In response to several B-cell activation signals.,PTM:Could be phosphorylated. Might be functionally regulated by protein kinase(s).,similarity:Contains 1 RGS domain.,tissue specificity:B-cell specific. Expression is relatively low in B-cells and chronic lymphocytic leukemia B-cells; however, in other types of malignant B-cell such as non-Hodgkin's lymphoma and

**Nanjing BYabscience technology Co.,Ltd**



hairy cell leukemia, expression is constitutively high.,

## Background

This gene encodes a member of the regulator of G-protein signalling family. This protein is located on the cytosolic side of the plasma membrane and contains a conserved, 120 amino acid motif called the RGS domain. The protein attenuates the signalling activity of G-proteins by binding to activated, GTP-bound G alpha subunits and acting as a GTPase activating protein (GAP), increasing the rate of conversion of the GTP to GDP. This hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. [provided by RefSeq, Jul 2008],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using RGS1 Monoclonal Antibody