



## Septin 14 Monoclonal Antibody

Catalog No     BYmab-04196       Isotype     IgG       Reactivity     Human;Rat;Mouse;       Applications     WB       Gene Name     14-9月       Protein Name     Septin-14       Immunogen     The antiserum was produced against synthesized peptide derived from human SEPT14. AA range:341-390       Specificity     Septin 14 Monoclonal Antibody detects endogenous levels of Septin 14 protein.       Formulation     Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.       Source     Monoclonal, Mouse, IgG       Purification     The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.       Dilution     WB 1:500-2000       Concentration     1 mg/ml       Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     SEPT14; Septin-14       Observed Band     50kD       Cell Pathway     Cytoplasm, Cytoplasm, cytoskeleton, Cell projection, axon, Cell projection, dendrite, Perikaryon, Cytoplasm, perinuclear region, Cytoplasmic vesicle, secretory vesics, acrossorial rese with projection is strys first. Published: 17922164), Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity).       Tissue Specificity     Testis-specific (at protein level).       Function     function:Involved in cytokinesis .,similarity:Belongs to the septin family, cytoskeletal proteins		
Reactivity         Human;Rat;Mouse;           Applications         WB           Gene Name         14-9 月           Protein Name         Septin-14           Immunogen         The antiserum was produced against synthesized peptide derived from human SEPT14. AA range:341-390           Specificity         Septin 14 Monoclonal Antibody detects endogenous levels of Septin 14 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         SEPT14; Septin-14           Observed Band         50kD           Cell Pathway         Cytoplasm. Cytoplasm, cytoplasm, perinuclear region. Cytoplasmic vesicle, secretory vesicle, acrosome. Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear im and manchette structure in early elongating spermatids during spermiogenesis (By similarity).           Tissue Specificity         Testis-specific (at protein level).           Function         function:Involved in cytokinesis .,similarity:Belongs to the septin family.	Catalog No	BYmab-04196
Applications WB  Gene Name 14-9月  Protein Name Septin-14  Immunogen The antiserum was produced against synthesized peptide derived from human SEPT14. AA range:341-390  Specificity Septin 14 Monoclonal Antibody detects endogenous levels of Septin 14 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms SEPT14; Septin-14  Observed Band 50kD  Cell Pathway Cytoplasm, Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear riegion . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity).  Tissue Specificity Testis-specific (at protein level).  Function function:Involved in cytokinesis .,similarity:Belongs to the septin family .,  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Isotype	IgG
Gene Name         14-9月           Protein Name         Septin-14           Immunogen         The antiserum was produced against synthesized peptide derived from human SEPT14. AA range:341-390           Specificity         Septin 14 Monoclonal Antibody detects endogenous levels of Septin 14 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dillution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         SEPT14; Septin-14           Observed Band         50kD           Cell Pathway         Cytoplasm Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear riegion . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)           Tissue Specificity         Testis-specific (at protein level).           Function         function:Involved in cytokinesis .,similarity:Belongs to the septin family . GTP-binding cytoskeletal proteins implicated in membrane transport, a	Reactivity	Human;Rat;Mouse;
Immunogen	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human SEPT14. AA range:341-390 Specificity Septin 14 Monoclonal Antibody detects endogenous levels of Septin 14 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year SEPT14; Septin-14 Observed Band Cell Pathway Cytoplasm. Cytoplasm, cytoskeleton. Cell projection, axon. Cell projection, dendrite. Perikaryon. Cytoplasm, perinuclear region. Cytoplasmic vesicle, secretory vesicle, acrosome. Colocalizes with actin reses fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity). Tissue Specificity Testis-specific (at protein level). Function SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular ferson et al.,	Gene Name	14-9月
SEPT14. AA range:341-390  Specificity Septin 14 Monoclonal Antibody detects endogenous levels of Septin 14 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms SEPT14; Septin-14  Observed Band 50kD  Cell Pathway Cytoplasm. Cytoplasm, cytoskeleton. Cell projection, axon. Cell projection, dendrite. Perikaryon. Cytoplasm, perinuclear region. Cytoplasmic vesicle, secretory vesicle, acrosome. Collocalizes with action stryss fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity).  Tissue Specificity Testis-specific (at protein level).  Function  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Protein Name	Septin-14
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms SEPT14; Septin-14  Observed Band 50kD  Cell Pathway Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed: 17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity).  Tissue Specificity Testis-specific (at protein level).  Function  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Immunogen	
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       SEPT14; Septin-14         Observed Band       50kD         Cell Pathway       Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)         Tissue Specificity       Testis-specific (at protein level).         Function       function:Involved in cytokinesis .,similarity:Belongs to the septin family ., cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al., cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al., cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Specificity	Septin 14 Monoclonal Antibody detects endogenous levels of Septin 14 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms SEPT14; Septin-14  Observed Band 50kD  Cell Pathway Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity).  Tissue Specificity Testis-specific (at protein level).  Function SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms SEPT14; Septin-14  Observed Band 50kD  Cell Pathway Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)  Tissue Specificity Testis-specific (at protein level).  Function function:Involved in cytokinesis .,similarity:Belongs to the septin family .,  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Source	Monoclonal, Mouse,lgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms SEPT14; Septin-14  Observed Band 50kD  Cell Pathway Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity).  Tissue Specificity Testis-specific (at protein level).  Function function:Involved in cytokinesis .,similarity:Belongs to the septin family.,  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms SEPT14; Septin-14  Observed Band 50kD  Cell Pathway Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)  Tissue Specificity Testis-specific (at protein level).  Function function:Involved in cytokinesis .,similarity:Belongs to the septin family.,  Background SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year  Synonyms  SEPT14; Septin-14  Observed Band  Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)  Tissue Specificity  Testis-specific (at protein level).  Function  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Concentration	1 mg/ml
Synonyms  SEPT14; Septin-14  Observed Band  Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity).  Tissue Specificity  Testis-specific (at protein level).  Function  function:Involved in cytokinesis .,similarity:Belongs to the septin family.,  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Purity	≥90%
Observed Band  Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)  Tissue Specificity  Testis-specific (at protein level).  Function  function:Involved in cytokinesis .,similarity:Belongs to the septin family.,  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm . Cytoplasm, cytoskeleton . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)  Tissue Specificity  Testis-specific (at protein level).  Function  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Synonyms	SEPT14; Septin-14
dendrite . Perikaryon . Cyfoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in early elongating spermatids during spermiogenesis (By similarity)  Tissue Specificity  Testis-specific (at protein level).  Function  function:Involved in cytokinesis .,similarity:Belongs to the septin family.,  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Observed Band	50kD
Function  function:Involved in cytokinesis .,similarity:Belongs to the septin family.,  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Cell Pathway	dendrite . Perikaryon . Cyfoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle, acrosome . Colocalizes with actin stress fibers (PubMed:17922164). Expressed in the perinuclear rim and manchette structure in
Background  SEPT14 is a member of the highly conserved septin family of GTP-binding cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Tissue Specificity	Testis-specific (at protein level).
cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,	Function	function:Involved in cytokinesis .,similarity:Belongs to the septin family.,
	Background	cytoskeletal proteins implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis, and other cellular functions (Peterson et al.,

Nanjing BYabscience technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using Septin 14 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd