



Septin 5 Monoclonal Antibody

Catalog No	BYmab-04199
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	5-9月
Protein Name	Septin-5
Immunogen	Synthesized peptide derived from Septin 5 . at AA range: 180-260
Specificity	Septin 5 Monoclonal Antibody detects endogenous levels of Septin 5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SEPT5; PNUTL1; Septin-5; Cell division control-related protein 1; CDCrel-1; Peanut-like protein 1
Observed Band	42kD
Cell Pathway	Cytoplasm . Cytoplasm, cytoskeleton . In platelets, found in areas surrounding alpha-granules.
Tissue Specificity	Expressed at high levels in the CNS, as well as in heart and platelets (at protein level).
Function	function:Involved in cytokinesis .,similarity:Belongs to the septin family.,subunit:May assemble into a multicomponent structure.,
Background	septin 5(SEPT5) Homo sapiens This gene is a member of the septin gene family of nucleotide binding proteins, originally described in yeast as cell division cycle regulatory proteins. Septins are highly conserved in yeast, Drosophila, and mouse and appear to regulate cytoskeletal organization. Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid cells. This gene is mapped to 22q11, the region frequently deleted in DiGeorge and velocardiofacial syndromes. A translocation involving the MLL gene and this gene has also been reported in patients with acute myeloid leukemia. Alternative

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务, 欢迎咨询

splicing results in multiple transcript variants. The presence of a non-consensus polyA signal (AACAAT) in this gene also results in read-through transcription into the downstream neighboring gene (GP1BB; platelet glycoprotein lb), whereby larger, non-coding transcripts are produced. [provided by RefSeq, Dec 2010], Avoid repeated freezing and thawing! matters needing attention This product can be used in immunological reaction related experiments. For Usage suggestions more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co., Ltd