



STEAP1 Monoclonal Antibody

| Catalog No BYmab-04227 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name STEAP1 Protein Name Metalloreductase STEAP1 Immunogen The antiserum was produced against synthesized peptide derived from human STEAP1. AA range;31-80 Specificity STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein . Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to re | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reactivity Human;Rat;Mouse; Applications WB Gene Name STEAP1 Protein Name Metalloreductase STEAP1 Immunogen The antiserum was produced against synthesized peptide derived from human STEAP1. AA range:31-80 Specificity STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed in prostate tumors. Function cofactor:FAD, function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor, similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxido | Catalog No | BYmab-04227 |
| Applications WB Gene Name STEAP1 Protein Name Metalloreductase STEAP1 Immunogen The antiserum was produced against synthesized peptide derived from human STEAP1. AA range:31-80 Specificity STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD. function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor. similarity:Belongs to the STEAP family. similarity:Contains 1 ferric oxidoreductase domain. tissue specificity. Ubiquitously expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Isotype | IgG |
| Gene Name STEAP1 Protein Name Metalloreductase STEAP1 Immunogen The antiserum was produced against synthesized peptide derived from human STEAP1. AA range:31-80 Specificity STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD, function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor. similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain, tissue specificity. Ubiquitously expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. | Reactivity | Human;Rat;Mouse; |
| Protein Name Metalloreductase STEAP1 Immunogen The antiserum was produced against synthesized peptide derived from human STEAP1. AA range:31-80 Specificity STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD., function: Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor., similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain., tissue specificity: Ubiquitously expressed. Highly expressed in prostate tumors. This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Applications | WB |
| Immunogen The antiserum was produced against synthesized peptide derived from human STEAP1. AA range:31-80 Specificity STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain, itssue specificity:Ubiquitously expressed in prostate tumors. Background This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a si | Gene Name | STEAP1 |
| STEAP1. AA range:31-80 Specificity STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD, function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain, tissue specificity: Ubiquitously expressed. Highly expressed in prostate tumors. Background This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Protein Name | Metalloreductase STEAP1 |
| Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD., function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor, similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain, tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Immunogen | |
| Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Celongs to the STEAP family, semilarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., Background This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Specificity | STEAP1 Monoclonal Antibody detects endogenous levels of STEAP1 protein. |
| Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane ; Multi-pass membrane protein . Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., Background This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD., function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor., similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain., tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors. Background This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Source | Monoclonal, Mouse,IgG |
| Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Purification | |
| Purity ≥90% Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., Background This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Dilution | WB 1:500-2000 |
| Storage Stability -20°C/1 year Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed in prostate tumors., Background This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Concentration | 1 mg/ml |
| Synonyms STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1 Observed Band 36kD Cell Pathway Endosome membrane; Multi-pass membrane protein. Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Purity | ≥90% |
| epithelial antigen of prostate 1 Observed Band Gell Pathway Endosome membrane; Multi-pass membrane protein. Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Storage Stability | -20°C/1 year |
| Cell Pathway Endosome membrane; Multi-pass membrane protein. Ubiquitously expressed. Highly expressed in prostate tumors. cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Synonyms | |
| Tissue Specificity Ubiquitously expressed. Highly expressed in prostate tumors. Cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Observed Band | 36kD |
| Function cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Cell Pathway | Endosome membrane ; Multi-pass membrane protein . |
| to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor, similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain, tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors., This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Tissue Specificity | Ubiquitously expressed. Highly expressed in prostate tumors. |
| upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen | Function | to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor, similarity:Belongs to the STEAP family, similarity:Contains 1 ferric oxidoreductase domain, tissue |
| | Background | upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen |

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using STEAP1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd