



TRAF2 Monoclonal Antibody

Catalog No	BYmab-04250
Isotype	IgG
Reactivity	Human;Rat;Mouse;;Bovine
Applications	WB
Gene Name	TRAF2
Protein Name	TNF receptor-associated factor 2
Immunogen	Synthesized peptide derived from the Internal region of human TRAF2.
Specificity	TRAF2 Monoclonal Antibody detects endogenous levels of TRAF2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TRAF2; TRAP3; TNF receptor-associated factor 2; E3 ubiquitin-protein ligase TRAF2; Tumor necrosis factor type 2 receptor-associated protein 3
Observed Band	60kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Brain,Cerebellum,Colon,Fetal brain,Human endometrium,Kidney,Leukocyte,Splee
Function	domain:The coiled coil domain mediates homo- and hetero-oligomerization.,domain:The MATH/TRAF domain binds to receptor cytoplasmic domains.,function:Adapter protein and signal transducer that links members of the tumor necrosis factor receptor family to different signaling pathways by association with the receptor cytoplasmic domain and kinases. Association to the receptor is also mediated by the interaction with TRADD. Mediates activation of NF-kappa-B and JNK and is involved in apoptosis. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. Seems to be involved in IL-15 signaling.,PTM:Ubiquitinated; mediated by SIAH2 and leading to its subsequent proteasomal degradation. Not ubiquitinated by SIAH1.,similarity:Contains 1 MATH domain.,similarity:Contains 1

Nanjing BYabscience technology Co.,Ltd



RING-type zinc finger.,similarity:Contains 2 TRAF-type zinc fingers.,subunit:Homotrimer (Pro

Background

TNF receptor associated factor 2(TRAF2) Homo sapiens The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kapMAB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can ubiquitinate and induce the degradation of this pro


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TRAF2 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd