



Wnt-1 Monoclonal Antibody

Catalog No BYmab-04269 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name WNT1 Protein Name Proto-oncogene Wnt-1 Immunogen The antiserum was produced against synthesized peptide derived from human WNT1. AA range;301-350 Specificity Wnt-1 Monocional Antibody detects endogenous levels of Wnt-1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monocional, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule impor		
Reactivity Applications WB Gene Name WNT1 Protein Name Proto-oncogene Wnt-1 Immunogen The antiserum was produced against synthesized peptide derived from human WNT1. AA range:301-350 Specificity Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters, similarity. Belongs to the Wnf family, subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3. The WNT gene family consists of structurally related genes which encode secreted signaling proleins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of tew WT gene family	Catalog No	BYmab-04269
Applications WB Gene Name WNT1 Protein Name Proto-oncogene Wnt-1 Immunogen The antiserum was produced against synthesized peptide derived from human WNT1. AA range:301-350 Specificity Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters, similarity. Belongs to the Wnt family., subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3. The WNT gene family consists of structurally related genes which encode secreted signaling proleins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of twWT gene family	Isotype	IgG
Gene Name Proto-oncogene Wnt-1 Immunogen The antiserum was produced against synthesized peptide derived from human WNT1. AA range:301-350 Specificity Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters, similarity Belongs to the Wnt family, subunit: Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Reactivity	Human;Mouse
Protein Name Proto-oncogene Wnt-1 Immunogen The antiserum was produced against synthesized peptide derived from human WNT1. AA range:301-350 Specificity Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable development. Is likely to signal over only few cell diameters, similarity:Belongs to the Wnt family, subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3. Background Secreted in Several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the Wnt gene family	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human WNT1. AA range:301-350 Specificity Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix. Secreted. Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable development. Is likely to signal over only few cell diameters, similarity. Belongs to the Wnt family, subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis.	Gene Name	WNT1
WNT1. AA range:301-350 Specificity Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix. Secreted. Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters., similarity: Belongs to the Wnt family., subunit: Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3. The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This sepens is a member of the WNT gene family	Protein Name	Proto-oncogene Wnt-1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix. Secreted. Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters, similarity:Belongs to the Wnt family.,subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters. similarity:Belongs to the Wnt family., subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3. Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Specificity	Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters. similarity: Belongs to the Wnt family., subunit: Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3. Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters., similarity:Belongs to the Wnt family., subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters.,similarity:Belongs to the Wnt family.,subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix . Secreted . Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters.,similarity:Belongs to the Wnt family.,subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix. Secreted. Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters., similarity: Belongs to the Wnt family., subunit: Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Concentration	1 mg/ml
Synonyms WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog Observed Band 45kD Cell Pathway Secreted, extracellular space, extracellular matrix. Secreted. Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters., similarity: Belongs to the Wnt family., subunit: Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3. Background The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family	Purity	≥90%
Observed Band Cell Pathway Secreted, extracellular space, extracellular matrix. Secreted. Tissue Specificity Testis, Function function: Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters., similarity: Belongs to the Wnt family., subunit: Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family.	Storage Stability	-20°C/1 year
Cell Pathway Secreted, extracellular space, extracellular matrix. Secreted. Tissue Specificity Testis, Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters.,similarity:Belongs to the Wnt family.,subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family.	Synonyms	WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog
Tissue Specificity Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters.,similarity:Belongs to the Wnt family.,subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family.	Observed Band	45kD
Function function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters., similarity:Belongs to the Wnt family., subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family.	Cell Pathway	Secreted, extracellular space, extracellular matrix . Secreted .
receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters., similarity:Belongs to the Wnt family., subunit:Interacts with PORCN. Interacts with RSPO1, RSPO2 and RSPO3., The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family.	Tissue Specificity	Testis,
secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family.	Function	receptors. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters.,similarity:Belongs to the Wnt family.,subunit:Interacts with PORCN.
	Background	secreted signaling proteins. These proteins have been implicated in oncogenesis

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



be 98% identical to the mouse Wnt1 protein at the amino acid	
mouse indicate that the Wnt1 protein functions in the induction	
mesencephalon and cerebellum. This gene was originally co	nsidered as a
candidate gene for Joubert syndrome, an autosomal recessi	ve disorder with
cerebellar hypoplasia as a leading feature. However, further	er studies suggested
that the gene mutations might not have a significant role in J	oubert syndrome.
This gene is clustered with another family member, WNT10E	B, in


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Wnt-1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd