



Elmo1 Monoclonal Antibody

Catalog No	BYmab-04297
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	ELMO1
Protein Name	Engulfment and cell motility protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from human Elmo1. AA range:22-71
Specificity	Elmo1 Monoclonal Antibody detects endogenous levels of Elmo1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ELMO1; KIAA0281; Engulfment and cell motility protein 1; Protein ced-12 homolog
Observed Band	83kD
Cell Pathway	Cytoplasm. Cell membrane. Translocation to plasma membrane seems to be mediated by DOCK1 and CRK.
Tissue Specificity	Widely expressed, with a higher expression in the spleen and placenta.
Function	function:Involved in cytoskeletal rearrangements required for phagocytosis of apoptotic cells and cell motility. Acts in association with DOCK1 and CRK. Was initially proposed to be required in complex with DOCK1 to activate Rac Rho small GTPases. May enhance the guanine nucleotide exchange factor (GEF) activity of DOCK1.,PTM:Phosphorylated by HCK.,similarity:Contains 1 ELMO domain.,similarity:Contains 1 PH domain.,subcellular location:Translocation to plasma membrane seems to be mediated by DOCK1 and CRK.,subunit:Interacts with BAI1 (By similarity). Interacts directly with the SH3-domain of DOCK1 via its SH3-binding site. Part of a complex with DOCK1 and RAC1. Part of a complex with DOCK1 and CRK isoform CRK-II. Interacts with PLEKHG6.,tissue

Nanjing BYabscience technology Co.,Ltd



specificity: Widely expressed, with a higher expression in the spleen and placenta.,

Background

This gene encodes a member of the engulfment and cell motility protein family. These proteins interact with dedicator of cytokinesis proteins to promote phagocytosis and cell migration. Increased expression of this gene and dedicator of cytokinesis 1 may promote glioma cell invasion, and single nucleotide polymorphisms in this gene may be associated with diabetic nephropathy. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Elmo1 Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658