



Macroglobulin α -2 Monoclonal Antibody

Catalog No	BYmab-04331
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	A2M
Protein Name	Alpha-2-macroglobulin
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human A2M. AA range:871-920
Specificity	Macroglobulin α -2 Monoclonal Antibody detects endogenous levels of Macroglobulin α -2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	$-20^{\circ}\text{C}/1$ year
Synonyms	A2M; CPAMD5; FWP007; Alpha-2-macroglobulin; Alpha-2-M; C3 and PZP-like alpha-2-macroglobulin domain-containing protein 5
Observed Band	170kD
Cell Pathway	Secreted .
Tissue Specificity	Secreted in plasma.
Function	developmental stage:Contrary to the rat protein, which is an acute phase protein, this protein is always present at high levels in circulation.,function:Is able to inhibit all four classes of proteinases by a unique 'trapping' mechanism. This protein has a peptide stretch, called the 'bait region' which contains specific cleavage sites for different proteinases. When a proteinase cleaves the bait region, a conformational change is induced in the protein which traps the proteinase. The entrapped enzyme remains active against low molecular weight substrates (activity against high molecular weight substrates is greatly reduced). Following cleavage in the bait region a thioester bond is hydrolyzed and mediates the covalent binding of the protein to the proteinase.,online information:Alpha-2 macroglobulin

Nanjing BYabscience technology Co.,Ltd



entry,similarity:Belongs to the protease inhibitor I39 (alpha-2-macroglobulin) family.,s

Background

Alpha-2-macroglobulin is a protease inhibitor and cytokine transporter. It inhibits many proteases, including trypsin, thrombin and collagenase. A2M is implicated in Alzheimer disease (AD) due to its ability to mediate the clearance and degradation of A-beta, the major component of beta-amyloid deposits. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using
Macroglobulin α -2 Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658