



SOCS-3 Monoclonal Antibody

Catalog No	BYmab-04354
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SOCS3 CIS3 SSI3
Protein Name	Suppressor of cytokine signaling 3 (SOCS-3) (Cytokine-inducible SH2 protein 3) (CIS-3) (STAT-induced STAT inhibitor 3) (SSI-3)
Immunogen	Synthetic peptide from human protein at AA range: 20-70
Specificity	The antibody detects endogenous SOCS-3
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Suppressor of cytokine signaling 3 (SOCS-3;Cytokine-inducible SH2 protein 3;CIS-3;STAT-induced STAT inhibitor 3;SSI-3)
Observed Band	25kD
Cell Pathway	intracellular,cytoplasm,cytosol,
Tissue Specificity	Widely expressed with high expression in heart, placenta, skeletal muscle, peripheral blood leukocytes, fetal and adult lung, and fetal liver and kidney. Lower levels in thymus.
Function	disease:Genetic variation in the promoter region of SOCS3 may be associated with susceptibility to atopic dermatitis 4 (ATOD4) [MIM:605805]. Atopic dermatitis [MIM:603165], also known as eczema commonly begins in infancy or early childhood and is characterized by ichy and inflamed skin.,domain:The ESS and SH2 domains are required for JAK phosphotyrosine binding. Further interaction with the KIR domain is necessary for signal and kinase inhibition.,domain:The SOCS box domain mediates the interaction with the Elongin BC complex, an adapter module in different E3 ubiquitin ligase complexes.,function:SOCS family proteins form part of a classical negative feedback system that regulates cytokine signal transduction. SOCS3 is involved in negative regulation of cytokines that

Nanjing BYabscience technology Co.,Ltd



signal through the JAK/STAT pathway. Inhibits cytokine signal transduction by binding to tyrosine kinase receptors includ

Background

This gene encodes a member of the STAT-induced STAT inhibitor (SSI), also known as suppressor of cytokine signaling (SOCS), family. SSI family members are cytokine-inducible negative regulators of cytokine signaling. The expression of this gene is induced by various cytokines, including IL6, IL10, and interferon (IFN)-gamma. The protein encoded by this gene can bind to JAK2 kinase, and inhibit the activity of JAK2 kinase. Studies of the mouse counterpart of this gene suggested the roles of this gene in the negative regulation of fetal liver hematopoiesis, and placental development. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658