



DAG1 Monoclonal Antibody

Catalog No	BYmab-04903
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DAG1
Protein Name	Dystroglycan (Dystrophin-associated glycoprotein 1) [Cleaved into: Alpha-dystroglycan (Alpha-DG); Beta-dystroglycan (Beta-DG)]
Immunogen	Synthesized peptide derived from human protein . at AA range: 830-910
Specificity	DAG1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	98kD
Cell Pathway	[Alpha-dystroglycan]: Secreted, extracellular space.; [Beta-dystroglycan]: Cell membrane ; Single-pass type I membrane protein. Cytoplasm, cytoskeleton. Nucleus, nucleoplasm . Cell membrane, sarcolemma . Cell junction, synapse, postsynaptic cell membrane . The monomeric form translocates to the nucleus via the action of importins and depends on RAN. Nuclear transport is inhibited by Tyr-892 phosphorylation. In skeletal muscle, this phosphorylated form locates to a vesicular internal membrane compartment. In muscle cells, sarcolemma localization requires the presence of ANK2, while localization to costameres requires the presence of ANK3. Localizes to neuromuscular junctions (NMJs) in the presence of ANK2 (By similarity). In peripheral nerves, localizes to the Schwann cell membrane. Colocal
Tissue Specificity	Expressed in a variety of fetal and adult tissues. In epidermal tissue, located to the basement membrane. Also expressed in keratinocytes and fibroblasts.
Function	function:Forms part of the dystrophin-associated protein complex (DAPC) which may link the cytoskeleton to the extracellular matrix. Alpha-dystroglycan functions

Nanjing BYabscience technology Co.,Ltd



as a laminin receptor. Binds to several types of arenaviruses. Is a target for the entry of Mycobacterium leprae into peripheral nerve Schwann cells.,online information:Dystroglycan entry,similarity:Contains 1 peptidase S72 domain.,subunit:Interacts with SGCD (By similarity). Interacts with AGR2 and AGR3.,tissue specificity:Expressed in a variety of fetal and adult tissues.,

Background

This gene encodes dystroglycan, a central component of dystrophin-glycoprotein complex that links the extracellular matrix and the cytoskeleton in the skeletal muscle. The encoded preproprotein undergoes O- and N-glycosylation, and proteolytic processing to generate alpha and beta subunits. Certain mutations in this gene are known to cause distinct forms of muscular dystrophy. Alternative splicing results in multiple transcript variants, all encoding the same protein. [provided by RefSeq, Nov 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using DAG1 Monoclonal Antibody