




LSM1 Monoclonal Antibody

Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity LSM1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye, Lymph node, Function function: Binds specifically to the 3'-terminal U-tract of U6 snRNA, similarity: Belongs to the snRNP Sm proteins family, subunit: LSm subunits form a heteromer with a donut shape, lissue specificity: Has elevated expression in pancreatic cancer and in several cancer-derived cell lines, Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-sidning proteins LSm protein stable heteromers that bind specifically to the 3-dapos; -terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/106 snRNA and may play a role in pre-mRNA splicing by mediating U4/106 snRNA and may play a role in rolellaring U4/106 in cellular in cellu		
Reactivity Human;Mouse Applications WB Gene Name LSM1 CASM Protein Name U6 snRNA-associated Sm-like protein LSm1 (Cancer-associated Sm-like) (Smanuclear ribonuclear CaSm) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity LSM1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye, Lymph node, Function function: Binds specifically to the 3'-terminal U-tract of U6 snRNA . similarity-Belongs to the snRNP Sm proteins family, subunit. Sm subunits form a heteromer with a donut shape, tissue specificity-Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated (LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3kapos;-terminal oligo(U) transformation and the progression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer derived in cellular transformation and the progression of several malignancies including lung cancer	Catalog No	BYmab-04953
Applications Gene Name LSM1 CASM Protein Name U6 snRNA-associated Sm-like protein LSm1 (Cancer-associated Sm-like) (Smanuclear ribonuclear CaSm) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity LSM1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye, Lymph node, function: Binds specifically to the 3'-terminal U-tract of U6 snRNA . similarity Belongs to the snRNP Sm proteins family . subunit; LSm subunits form a heteromer with a donut shape, tissue specificity/Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated (LSM1) Homo sapiers This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3'sapos;-terminal oliqu(U) transformation and the progression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer transformation and the progression of several malignancies including lung cancer.	Isotype	IgG
Gene Name LSM1 CASM Protein Name U6 snRNA-associated Sm-like protein LSm1 (Cancer-associated Sm-like) (Smanuclear ribonuclear CaSm) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity LSM1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.; similarity. Belongs to the snRNP Sm proteins family, subunit:LSm subunits form a heteromer with a donut shape, dissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) trace of U6 snRNA and may play a ro	Reactivity	Human;Mouse
Protein Name U6 snRNA-associated Sm-like protein LSm1 (Cancer-associated Sm-like) (Smanuclear ribonuclear CaSm) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity LSM1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, function:Binds specifically to the 3'-terminal U-tract of U6 snRNA., similarity. Belongs to the snRNP Sm proteins family, subunit:LSm subunits form a heteromer with a donut shape, dissue specificity. Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) to 10 snRNP after proteirs of lein pre-mRNA splicing by mediating U4/106 snRNP formation. Increased expression of this gene may play a role in relular transformation and the proquession of several malignancies including lung cancer.	Applications	WB
Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity LSM1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye, Lymph node, Function function: Binds specifically to the 3'-terminal U-tract of U6 snRNA, similarity: Belongs to the snRNP Sm proteins family, subunit: LSm subunits form a heteromer with a donut shape, tissue specificity: Has elevated expression in pancreatic cancer and in several cancer-derived cell lines, but of U6 snRNA and may play a role in pre-miXNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellulating transformation and the progression of several mail(anganches including lung cancer	Gene Name	LSM1 CASM
Specificity LSM1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye, Lymph node, Function function: Binds specifically to the 3'-terminal U-tract of U6 snRNA., similarity: Belongs to the snRNP Sm proteins family., subunit: LSm subunits form a heteromer with a donut shape, tissue specificity: Has elevated expression in pancreatic cancer and in several cancer-derived cell lines. Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3' terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of several malignancies including lung cancer	Protein Name	U6 snRNA-associated Sm-like protein LSm1 (Cancer-associated Sm-like) (Small nuclear ribonuclear CaSm)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye, Lymph node, Function function: Binds specifically to the 3'-terminal U-tract of U6 snRNA. similarity: Belongs to the snRNP Sm proteins family, subunit: LSm subunits form a heteromer with a donut shape. tissue specificity: Has elevated expression in pancreatic cancer and in several cancer-derived cell lines. Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3' terminal oligo(U) trace of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer	Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, Function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA_similarity:Belongs to the snRNP Sm proteins family, subunit:LSm subunits form a heteromer with a donut shape, tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines. Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) trace of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer	Specificity	LSM1 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, Function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family,.subunit:LSm subunits form a heteromer with a donut shape, tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of several malignancies including lung cancer.	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, function: Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3'-terminal oligo(U) traof U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, Function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3'-terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer.	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, Function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3'-terminal oligo(U) trace of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer.	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, Function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer.	Concentration	1 mg/ml
Synonyms Observed Band 14kD Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, function: Binds specifically to the 3'-terminal U-tract of U6 snRNA., similarity: Belongs to the snRNP Sm proteins family., subunit: LSm subunits form a heteromer with a donut shape., tissue specificity: Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm protein form stable heteromers that bind specifically to the 3'-terminal oligo(U) trace of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer.	Purity	≥90%
Cell Pathway Cytoplasm . Cytoplasm, P-body . Tissue Specificity Eye,Lymph node, function:Binds specifically to the 3'-terminal U-tract of U6 snRNA., similarity:Belongs to the snRNP Sm proteins family., subunit:LSm subunits form a heteromer with a donut shape., tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) traced of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer.	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Cytoplasm, P-body . Eye,Lymph node, function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer	Synonyms	
Tissue Specificity Eye,Lymph node, function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer	Observed Band	14kD
Function function:Binds specifically to the 3'-terminal U-tract of U6 snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer	Cell Pathway	Cytoplasm . Cytoplasm, P-body .
snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated expression in pancreatic cancer and in several cancer-derived cell lines., Background LSM1 homolog, mRNA degradation associated(LSM1) Homo sapiens This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) tra of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer.	Tissue Specificity	Eye,Lymph node,
gene encodes a member of the LSm family of RNA-binding proteins. LSm proteir form stable heteromers that bind specifically to the 3'-terminal oligo(U) trate of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cance	Function	snRNA.,similarity:Belongs to the snRNP Sm proteins family.,subunit:LSm subunits form a heteromer with a donut shape.,tissue specificity:Has elevated
	Background	gene encodes a member of the LSm family of RNA-binding proteins. LSm proteins form stable heteromers that bind specifically to the 3'-terminal oligo(U) tract of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Increased expression of this gene may play a role in cellular transformation and the progression of several malignancies including lung cancer,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








	been observed for this gene, and a pseudogene of this gene is located on the short arm of chromosome 9. [provided by RefSeq, Nov 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658