



## FHL3 Monoclonal Antibody

Immunogen   Synthesized peptide derived from human protein . at AA range: 40-120		
Reactivity Human;Mouse  Applications WB  Gene Name FHL3 SLIM2  Protein Name Four and a half LIM domains protein 3 (FHL-3) (Skeletal muscle LIM-protein (SLIM-2)  Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity FHL3 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains, tissue specificity:Expressed only in skeletal muscle.  Background The protein encoded by this gene is a member of a family of proteins contented by motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle requested muscles a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Catalog No	BYmab-04989
Applications  WB  Gene Name FHL3 SLIM2  Protein Name Four and a half LIM domains protein 3 (FHL-3) (Skeletal muscle LIM-protein (SLIM-2)) Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity FHL3 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity:Contains 3 LIM zinc-binding domains, similarity:Contains 4 LIM zinc-binding domains, tissue specificity:Expressed only in skeletal muscle.  Background The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental requiators SMAD2, SMAD3, and SMAD4, the skeletal muscle.	Isotype	IgG
Gene Name FHL3 SLIM2  Protein Name Four and a half LIM domains protein 3 (FHL-3) (Skeletal muscle LIM-protein (SLIM-2)  Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity FHL3 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity:Contains 3 LIM zinc-binding domains, similarity:Contains 4 LIM zinc-binding domains, tissue specificity:Expressed only in skeletal muscle.  Background The encoded by this gene is a member of a family of proteins conductive developmental requiators SMAD2, SMAD3, and SMAD3, and SMAD4, the skeletal muscle developmental requiators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Reactivity	Human;Mouse
Protein Name   Four and a half LIM domains protein 3 (FHL-3) (Skeletal muscle LIM-protein (SLIM-2)   Immunogen   Synthesized peptide derived from human protein . at AA range: 40-120   Specificity   FHL3 Monoclonal Antibody detects endogenous levels of protein.   Formulation   Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source   Monoclonal, Mouse, IgG   Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution   WB 1:500-2000   Concentration   1 mg/ml   Purity   ≥90%   Storage Stability   -20°C/1 year   Synonyms   Observed Band   30kD   Cell Pathway   Nucleus . Cytoplasm .   Tissue Specificity   Expressed only in skeletal muscle.   Function   similarity: Contains 3 LIM zinc-binding domains, similarity: Contains 4 LIM zinc-binding domains, tissue specificity: Expressed only in skeletal muscle.   Background   The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finget motif. The encoded protein has been shown to interact with the cancer developmental requiators SMAD2, SMAD3, and SMAD4, the skeletal muscle.	Applications	WB
Immunogen   Synthesized peptide derived from human protein . at AA range: 40-120	Gene Name	FHL3 SLIM2
Specificity FHL3 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, lgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity: Contains 3 LIM zinc-binding domains., similarity: Contains 4 LIM zinc-binding domains., tissue specificity: Expressed only in skeletal muscle.,  Background The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, the skeletal muscle.	Protein Name	Four and a half LIM domains protein 3 (FHL-3) (Skeletal muscle LIM-protein 2) (SLIM-2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  Background The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD4, she skeletal muscle.	Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       30kD         Cell Pathway       Nucleus . Cytoplasm .         Tissue Specificity       Expressed only in skeletal muscle.         Function       similarity: Contains 3 LIM zinc-binding domains., similarity: Contains 4 LIM zinc-binding domains., tissue specificity: Expressed only in skeletal muscle.,         Background       The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Specificity	FHL3 Monoclonal Antibody detects endogenous levels of protein.
Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band       30kD         Cell Pathway       Nucleus . Cytoplasm .         Tissue Specificity       Expressed only in skeletal muscle.         Function       similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,         Background       The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  Background The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  Background The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  Background The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle.	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year  Synonyms  Observed Band 30kD  Cell Pathway Nucleus . Cytoplasm .  Tissue Specificity Expressed only in skeletal muscle.  Function similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  Background The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Concentration	1 mg/ml
Synonyms  Observed Band  Observed Band  Cell Pathway  Nucleus . Cytoplasm .  Tissue Specificity  Expressed only in skeletal muscle.  Function  similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Purity	≥90%
Observed Band  Cell Pathway  Nucleus . Cytoplasm .  Expressed only in skeletal muscle.  Function  similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Storage Stability	-20°C/1 year
Cell Pathway       Nucleus . Cytoplasm .         Tissue Specificity       Expressed only in skeletal muscle.         Function       similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,         Background       The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Synonyms	
Tissue Specificity  Expressed only in skeletal muscle.  Function  similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle		
Function  similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Observed Band	30kD
Zinc-binding domains.,tissue specificity:Expressed only in skeletal muscle.,  The protein encoded by this gene is a member of a family of proteins conta a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle		
a four-and-a-half LIM domain, which is a highly conserved double zinc finger motif. The encoded protein has been shown to interact with the cancer developmental regulators SMAD2, SMAD3, and SMAD4, the skeletal muscle	Cell Pathway	Nucleus . Cytoplasm .
myogenesis protein MyoD, and the high-affinity IgE beta chain regulator MZI This protein may be involved in tumor suppression, repression of MyoD expression, and repression of IgE receptor expression. Two transcript variar encoding different isoforms have been found for this gene. [provided by Refs Aug 2011],	Cell Pathway Tissue Specificity	Nucleus . Cytoplasm .  Expressed only in skeletal muscle.  similarity:Contains 3 LIM zinc-binding domains.,similarity:Contains 4 LIM

Nanjing BYabscience technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using FHL3 Monoclonal Antibody