





MTF1 Monoclonal Antibody

Catalog No BYmab-04990 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name MTF1 Protein Name Metal regulatory transcription factor 1 (MRE-binding transcription factor) (Transcription factor MTF-1) Immunogen Synthesized peptide derived from human protein . at AA range: 260-340 Specificity MTF1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals . Tissue Specificity Liver,Muscle, Function function:Activates the metallothionein I promoter. Binds to the metal responsive element (MRE), ismilarity.Contains 6 CZH2-type zinc fingers. Background		
Applications WB Gene Name MTF1 Protein Name Metal regulatory transcription factor 1 (MRE-binding transcription factor) (Transcription factor MTF-1) Immunogen Synthesized peptide derived from human protein . at AA range: 260-340 Specificity MTF1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver,Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE), similarity. Contains 6 C2H2-type zinc fingers., This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], Matters needing Avoid repeated freezing and thawing!	Catalog No	BYmab-04990
Applications WB Gene Name MTF1 Protein Name (Iranscription factor MTF-1) Immunogen Synthesized peptide derived from human protein . at AA range: 260-340 Specificity MTF1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals . Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE)., similarity. Contains 6 C2H2-type zinc fingers., This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shutting protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], Matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name MTF1 Protein Name Metal regulatory transcription factor 1 (MRE-binding transcription factor) (Transcription factor MTF-1) Immunogen Synthesized peptide derived from human protein . at AA range: 260-340 Specificity MTF1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals . Tissue Specificity Liver,Muscle, Function function:Activates the metallothionein I promoter. Binds to the metal responsive element (MRE), similarity:Contains 6 C2H2-type zinc fingers. Background This gene encodes a transcription factor that induces expression of metallotinoneins and other genes involved in metal homeostasis in response to heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq,	Reactivity	Human;Mouse
Protein Name (Metal regulatory transcription factor 1 (MRE-binding transcription factor) (Transcription factor MTF-1) Immunogen Synthesized peptide derived from human protein . at AA range: 260-340 Specificity MTF1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE), similarity: Contains 6 C2H2-type zinc fingers., This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Applications	WB
Immunogen Synthesized peptide derived from human protein . at AA range: 260-340 Specificity MTF1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals . Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE), similarity: Contains 6 C2H2-type zinc fingers. Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing	Gene Name	MTF1
Specificity MTF1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein 1 promoter. Binds to the metal responsive element (MRE) similarity: Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Protein Name	Metal regulatory transcription factor 1 (MRE-binding transcription factor) (Transcription factor MTF-1)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein 1 promoter. Binds to the metal responsive element (MRE), similarity: Contains 6 C2H2-type zinc fingers., This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human protein . at AA range: 260-340
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals . Tissue Specificity Liver,Muscle, Function function:Activates the metallothionein I promoter. Binds to the metal responsive element (MRE), similarity:Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Specificity	MTF1 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver,Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE).,similarity: Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals . Tissue Specificity Liver,Muscle, Function function:Activates the metallothionein I promoter. Binds to the metal responsive element (MRE).,similarity:Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein 1 promoter. Binds to the metal responsive element (MRE)., similarity: Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver,Muscle, Function function:Activates the metallothionein I promoter. Binds to the metal responsive element (MRE).,similarity:Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Synonyms Observed Band Each Specificity Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE)., similarity: Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 82kD Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE)., similarity: Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, Function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE)., similarity: Contains 6 C2H2-type zinc fingers., Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals Tissue Specificity Liver, Muscle, function function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE)., similarity: Contains 6 C2H2-type zinc fingers., This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Liver,Muscle, function function:Activates the metallothionein I promoter. Binds to the metal responsive element (MRE).,similarity:Contains 6 C2H2-type zinc fingers., This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Observed Band	82kD
Function function:Activates the metallothionein I promoter. Binds to the metal responsive element (MRE).,similarity:Contains 6 C2H2-type zinc fingers., This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Cell Pathway	Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals
Background This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Liver, Muscle,
metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Function	function: Activates the metallothionein I promoter. Binds to the metal responsive element (MRE)., similarity: Contains 6 C2H2-type zinc fingers.,
	Background	metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element
		Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd