



LMTK1 Monoclonal Antibody

| Catalog No | BYmab-05002 |
|--|--|
| Isotype | lgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | AATK AATYK KIAA0641 LMR1 LMTK1 |
| Protein Name | Serine/threonine-protein kinase LMTK1 (EC 2.7.11.1) (Apoptosis-associated tyrosine kinase) (AATYK) (Brain apoptosis-associated tyrosine kinase) (CDK5-binding protein) (Lemur tyrosine kinase 1) (p35-bi |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 1110-1190 |
| Specificity | LMTK1 Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| | |
| Concentration | 1 mg/ml |
| Concentration Purity | 1 mg/ml ≥90% |
| | |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% |
| Purity Storage Stability Synonyms | ≥90% -20°C/1 year |
| Purity Storage Stability Synonyms Observed Band | ≥90% -20°C/1 year 151kD Membrane ; Single-pass type I membrane protein . Cytoplasm . Cytoplasm, |
| Purity Storage Stability Synonyms Observed Band Cell Pathway | ≥90% -20°C/1 year 151kD Membrane ; Single-pass type I membrane protein . Cytoplasm . Cytoplasm, perinuclear region . Predominantly perinuclear. |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | ≥90% -20°C/1 year 151kD Membrane ; Single-pass type I membrane protein . Cytoplasm . Cytoplasm, perinuclear region . Predominantly perinuclear. Expressed in brain. catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:May be involved in neuronal differentiation.,induction:Up-regulated during apoptosis.,PTM:Autophosphorylated. Phosphorylated by CDK5.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Contains 1 protein kinase domain.,subcellular location:Predominantly perinuclear.,subunit:Interacts with CDK5.,tissue |



国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务,欢迎咨询

| | for the induction of growth arrest and/or apoptosis of myeloid precursor cells. This gene has been shown to produce neuronal differentiation in a neuroblastoma cell line. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2011], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| Products Images | |

Nanjing BYabscience technology Co.,Ltd