



KAT6B Monoclonal Antibody

Catalog No	BYmab-05011
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	KAT6B KIAA0383 MORF MOZ2 MYST4
Protein Name	Histone acetyltransferase KAT6B (EC 2.3.1.48) (Histone acetyltransferase MOZ2) (MOZ, YBF2/SAS3, SAS2 and TIP60 protein 4) (MYST-4) (Monocytic leukemia zinc finger protein-related factor)
Immunogen	Synthesized peptide derived from human protein . at AA range: 1170-1250
Specificity	KAT6B Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	228kD
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitously expressed, with high levels in heart, pancreas, testis and ovary.
Function	catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,disease:A chromosomal aberration involving MYST4 may be a cause acute myeloid leukemias. Translocation t(10;16)(q22;p13) with CREBBP.,domain:The N-terminus is involved in transcriptional activation while the C-terminus is involved in transcriptional repression.,function:Histone acetyltransferase which may be involved in both positive and negative regulation of transcription. Required for RUNX2-dependent transcriptional activation. May be involved in cerebral cortex development. Component of the MOZ/MORF complex which has a histone H3 acetyltransferase activity.,PTM:Autoacetylated.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the MYST (SAS/MOZ) family.,similarity:Contains 1 C2HC-type zinc finger.,similarity:Contains 2

Nanjing BYabscience technology Co.,Ltd



PHD-type zinc fingers.,subunit:Component of the MOZ/MORF composed

Background

The protein encoded by this gene is a histone acetyltransferase and component of the MOZ/MORF protein complex. In addition to its acetyltransferase activity, the encoded protein has transcriptional activation activity in its N-terminal end and transcriptional repression activity in its C-terminal end. This protein is necessary for RUNX2-dependent transcriptional activation and could be involved in brain development. Mutations have been found in patients with genitopatellar syndrome. A translocation of this gene and the CREBBP gene results in acute myeloid leukemias. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2012],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658