



HJURP Monoclonal Antibody

Catalog No	BYmab-05070
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	HJURP FAKTS FLEG1 URLC9
Protein Name	Holliday junction recognition protein (14-3-3-associated AKT substrate) (Fetal liver-expressing gene 1 protein) (Up-regulated in lung cancer 9)
Immunogen	Synthesized peptide derived from human protein . at AA range: 80-160
Specificity	HJURP Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	82kD
Cell Pathway	Nucleus, nucleolus. Chromosome, centromere. Localizes in centromeres during late telophase and early G1, when CENPA nucleosomes are assembled. Localizes to nucleolus during S phase, nucleolus site being often related to storage.
Tissue Specificity	According to PubMed:17256767, highly expressed in the thymus with lower levels in the placenta, small intestine, liver, skeletal muscle, and colon. According to PubMed:17823411, highly expressed in testis, and at a relatively lower level in thymus and bone marrow. Significantly overexpressed in many lung cancer samples, compared with normal lung.
Function	function:Centromeric protein that plays a central role in the incorporation and maintenance of histone H3-like variant CENPA at centromeres. Acts as a specific chaperone for CENPA and is required for the incorporation of newly synthesized CENPA molecules into nucleosomes at replicated centromeres. Directly binds Holliday junctions.,sequence caution:Translated as Arg.,subcellular location:Localizes in centromeres during late telophase and early G1, when

Nanjing BYabscience technology Co.,Ltd



	<p>CENPA nucleosomes are assembled. Localizes to nucleolus during S phase, nucleolus site being often related to storage.,subunit:Interacts with CENPA (via CATD domain); the interaction is direct and specific for CENPA since it does not interact with H3.1- or H3.3-containing nucleosomes. Interacts with 14-3-3 family members in a phosphorylation-dependent manner. Interacts with MSH5 and NBN.,tissue specificity:According to PubMed:17256767 high</p>
Background	<p>function:Centromeric protein that plays a central role in the incorporation and maintenance of histone H3-like variant CENPA at centromeres. Acts as a specific chaperone for CENPA and is required for the incorporation of newly synthesized CENPA molecules into nucleosomes at replicated centromeres. Directly binds Holliday junctions.,sequence caution:Translated as Arg.,subcellular location:Localizes in centromeres during late telophase and early G1, when CENPA nucleosomes are assembled. Localizes to nucleolus during S phase, nucleolus site being often related to storage.,subunit:Interacts with CENPA (via CATD domain); the interaction is direct and specific for CENPA since it does not interact with H3.1- or H3.3-containing nucleosomes. Interacts with 14-3-3 family members in a phosphorylation-dependent manner. Interacts with MSH5 and NBN.,tissue specificity:According to PubMed:17256767 highly expressed in the thymus with lower levels in the placenta, small intestine, liver, skeletal muscle, and colon. According to PubMed:17823411 highly expressed in testis, and at a relatively lower level in thymus and bone marrow. Significantly overexpressed in many lung cancer samples, compared with normal lung.,</p>
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images