



GOT1B Monoclonal Antibody


| Catalog No | BYmab-05194 |
|-----------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | GOLT1B GOT1A CGI-141 HDCMA39P UNQ432/PRO793 |
| Protein Name | Vesicle transport protein GOT1B (Golgi transport 1 homolog B) (Putative NF-kappa-B-activating protein 470) (hGOT1a) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 70-150 |
| Specificity | GOT1B Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| | 15kD |
| Cell Pathway | 15kD Golgi apparatus membrane ; Multi-pass membrane protein . |
| Cell Pathway Tissue Specificity | |
| _ | Golgi apparatus membrane ; Multi-pass membrane protein . Widely expressed. Tends to be up-regulated in seminomas compared to normal |
| Tissue Specificity | Golgi apparatus membrane ; Multi-pass membrane protein . Widely expressed. Tends to be up-regulated in seminomas compared to normal testis. function:May be involved in fusion of ER-derived transport vesicles with the Golgi |
| Tissue Specificity Function | Golgi apparatus membrane; Multi-pass membrane protein. Widely expressed. Tends to be up-regulated in seminomas compared to normal testis. function:May be involved in fusion of ER-derived transport vesicles with the Golgi complex.,similarity:Belongs to the GOT1 family., function:May be involved in fusion of ER-derived transport vesicles with the Golgi |
| Tissue Specificity Function Background matters needing | Golgi apparatus membrane; Multi-pass membrane protein. Widely expressed. Tends to be up-regulated in seminomas compared to normal testis. function:May be involved in fusion of ER-derived transport vesicles with the Golgi complex.,similarity:Belongs to the GOT1 family., function:May be involved in fusion of ER-derived transport vesicles with the Golgi complex.,similarity:Belongs to the GOT1 family., |

Nanjing BYabscience technology Co.,Ltd





Products Images



Western Blot analysis of various cells using GOT1B Monoclonal Antibody

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658