



# PER1 Monoclonal Antibody

<b>Catalog No</b>	BYmab-05202
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	PER1 KIAA0482 PER RIGUI
<b>Protein Name</b>	Period circadian protein homolog 1 (hPER1) (Circadian clock protein PERIOD 1) (Circadian pacemaker protein Rigiui)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 1110-1190
<b>Specificity</b>	PER1 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	141kD
<b>Cell Pathway</b>	Nucleus. Cytoplasm. Nucleocytoplasmic shuttling is effected by interaction with other circadian core oscillator proteins and/or by phosphorylation. Retention of PER1 in the cytoplasm occurs through PER1-PER2 heterodimer formation. Translocate to the nucleus after phosphorylation by CSNK1D or CSNK1E. Also translocated to the nucleus by CRY1 or CRY2 (By similarity). .
<b>Tissue Specificity</b>	Widely expressed. Expressed in hair follicles (at protein level).Found in heart, brain, placenta, lung, liver, skeletal muscle, pancreas, kidney, spleen, thymus, prostate, testis, ovary and small intestine. Highest level in skeletal muscle.
<b>Function</b>	alternative products:Additional isoforms seem to exist,function:Component of the circadian clock mechanism which is essential for generating circadian rhythms. Negative element in the circadian transcriptional loop. Influences clock function by interacting with other circadian regulatory proteins and transporting them to the nucleus. Negatively regulates CLOCK NPAS2-BMAL1 BMAL2-induced transactivation.,induction:Serum-induced levels in fibroblasts show circadian oscillations. Maximum levels after 1 hour stimulation, minimum levels after 12

**Nanjing BYabscience technology Co.,Ltd**



hours. Another peak is then observed after 20 hours. Protein levels show maximum levels at 6 hours, decrease to reach minimum levels at 20 hours, and increase again to reach a second peak after 26 hours. Levels then decrease slightly and then increase to maximum levels at 32 hours. Levels of phosphorylated form increase between 3 hours and 12 hours.,PT

**Background**

This gene is a member of the Period family of genes and is expressed in a circadian pattern in the suprachiasmatic nucleus, the primary circadian pacemaker in the mammalian brain. Genes in this family encode components of the circadian rhythms of locomotor activity, metabolism, and behavior. This gene is upregulated by CLOCK/ARNTL heterodimers but then represses this upregulation in a feedback loop using PER/CRY heterodimers to interact with CLOCK/ARNTL. Polymorphisms in this gene may increase the risk of getting certain cancers. Alternative splicing has been observed in this gene; however, these variants have not been fully described. [provided by RefSeq, Jan 2014],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

**Nanjing BYabscience technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658