



RM19 Monoclonal Antibody

Catalog NoBYmab-05226isotypeIgGReactivityHuman;MouseApplicationsWBGene NameMRPL19 KIAA0104 MRPL15Protein Name39S ribosomal protein L19, mitochondrial (L19mt) (MRP-L19) (39S ribos protein L15, mitochondrial) (L15mt) (MRP-L15)ImmunogenSynthesized peptide derived from human protein . at AA range: 160-240SpecificityRM19 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSynondrion .Tissue SpecificityB-cell.Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal protein L19P family.,	
ReactivityHuman;MouseApplicationsWBGene NameMRPL19 KIAA0104 MRPL15Protein Name39S ribosomal protein L19, mitochondrial (L19mt) (MRP-L19) (39S ribos protein L15, mitochondrial) (L15mt) (MRP-L19) (39S ribos protein L15, mitochondrial) (L15mt) (MRP-L19)ImmunogenSynthesized peptide derived from human protein . at AA range: 160-240SpecificityRM19 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsUitochondrion .Cell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
ApplicationsWBGene NameMRPL19 KIAA0104 MRPL15Protein Name39S ribosomal protein L19, mitochondrial (L19mt) (MRP-L19) (39S ribosomal protein L15, mitochondrial) (L19mt) (MRP-L15)ImmunogenSynthesized peptide derived from human protein . at AA range: 160-240SpecificityRM19 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsUitochondrion .Charles SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g mitopeomes, where this ratio is reversed. Another difference between marrow	
Gene Name MRPL19 KIAA0104 MRPL15 Protein Name 39S ribosomal protein L19, mitochondrial (L19mt) (MRP-L19) (39S ribos protein L15, mitochondrial) (L15mt) (MRP-L15) Immunogen Synthesized peptide derived from human protein . at AA range: 160-240 Specificity RM19 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Mitochondrion . Tissue Specificity B-cell,Bone marrow, Function similarity:Belongs to the ribosomal protein L19P family., Background Mammalian mitochondrial ribosomal protein L19P family.,	
Protein Name 39S ribosomal protein L19, mitochondrial (L19mt) (MRP-L19) (39S ribos protein L15, mitochondrial) (L15mt) (MRP-L15) Immunogen Synthesized peptide derived from human protein . at AA range: 160-240 Specificity RM19 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20° C/1 year Synonyms Observed Band 32kD Cell Pathway Mitochondrion . Tissue Specificity B-cell,Bone marrow, Function similarity:Belongs to the ribosomal protein L19P family., Background Mammalian mitochondrial ribosomal protein L19P family.,	
ImmunogenSynthesized peptide derived from human protein . at AA range: 160-240SpecificityRM19 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms32kDCell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal protein sare encoded by nuclear g ney hep in protein synthesis within the mitochondrial ribosomal protein sare encoded by nuclear g ney hep in protein synthesis within the mitochondrial ribosomal protein between mar ar abos compact of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ar bio serves A. Another difference between mar	
SpecificityRM19 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsImage: StabilityObserved Band32kDCell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal protein Serven difference between marriers within the mitochondrial ribosomal protein Compared to proka mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka were this ratio is reversed. Another difference between marriers are seried between ma	somal
FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsObserved BandCell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrial ribosomal arbiosomes) consist of a small 28S subunit and a large 39S subunit, have an estimated 75% protein to rRNA composition compared to proka arbiosomes) consist of a small 28S subunit and a large 39S subunit, have an estimated 75% protein to rRNA composition compared to proka arbiosomes) consist of a small 28S subunit and a large 39S subunit, have an estimated 75% protein to rRNA composition compared to proka arbiosomes) consist of a small 28S subunit and a large 39S subunit, have an estimated 75% protein to rRNA composition compared to proka arbiosomes) consist of a small 28S subunit and a large 39S subunit, have an estimated 75% protein to rRNA composition compared to proka arbiosomes) consist of a small 28S subunit and a large 39S subunit, have an estimated 75% protein to rRNA composition compared to proka arbiosomes, where this ratio is reversed. Another difference between mar)
SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsUnderstandObserved Band32kDCell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 32kD Cell Pathway Mitochondrion . Tissue Specificity B-cell,Bone marrow, Function similarity:Belongs to the ribosomal protein L19P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion compared to proka ribosomes, where this ratio is reversed. Another difference between marribosomes, where this ratio is reversed. Another difference between marribosomes.	
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 32kD Cell Pathway Mitochondrion . Tissue Specificity B-cell,Bone marrow, Function similarity:Belongs to the ribosomal protein L19P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosor (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 32kD Cell Pathway Mitochondrion . Tissue Specificity B-cell,Bone marrow, Function similarity:Belongs to the ribosomal protein L19P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between marrow	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 32kD Cell Pathway Mitochondrion . Tissue Specificity B-cell,Bone marrow, Function similarity:Belongs to the ribosomal protein L19P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosomes, where this ratio is reversed. Another difference between marrow	
Storage Stability-20°C/1 yearSynonyms-20°C/1 yearObserved Band32kDCell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrial ribosom ribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
SynonymsObserved Band32kDCell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
Observed Band32kDCell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mark	
Cell PathwayMitochondrion .Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mark	
Tissue SpecificityB-cell,Bone marrow,Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
Functionsimilarity:Belongs to the ribosomal protein L19P family.,BackgroundMammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear g help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
help in protein synthesis within the mitochondrion. Mitochondrial ribosom (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. have an estimated 75% protein to rRNA composition compared to proka ribosomes, where this ratio is reversed. Another difference between mar	
mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S n Among different species, the proteins comprising the mitoribosome differ in sequence, and sometimes in biochemical properties, which prevents e recognition by sequence homology. This gene encodes a 39S subunit pr	nes They aryotic mmalian rRNA. r greatly easy

Nanjing BYabscience technology Co.,Ltd





Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
matters needing attention	Avoid repeated freezing and thawing!
	[provided by RefSeq, Jul 2008],

Products Images



Nanjing BYabscience technology Co.,Ltd