



3HAO Monoclonal Antibody

Catalog No	BYmab-05236
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HAAO
Protein Name	3-hydroxyanthranilate 3,4-dioxygenase (EC 1.13.11.6) (3-hydroxyanthranilate oxygenase) (3-HAO) (3-hydroxyanthranilic acid dioxygenase) (HAD)
Immunogen	Synthesized peptide derived from human protein . at AA range: 30-110
Specificity	3HAO Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	31kD
Cell Pathway	Cytoplasm, cytosol .
Tissue Specificity	Brain,Hippocampus,Placenta,
Function	catalytic activity:3-hydroxyanthranilate + O(2) = 2-amino-3-carboxymuconate semialdehyde.,cofactor:Fe(2+) ion.,function:Catalyzes the oxidative ring opening of 3-hydroxyanthranilate to 2-amino-3-carboxymuconate semialdehyde, which spontaneously cyclizes to quinolinate.,pathway:Cofactor biosynthesis; NAD(+) biosynthesis; pyridine-2,3-dicarboxylate from L-kynurenine: step 3/3.,similarity:Belongs to the 3-HAO family.,subunit:Monomer.,
Background	3-Hydroxyanthranilate 3,4-dioxygenase is a monomeric cytosolic protein belonging to the family of intramolecular dioxygenases containing nonheme ferrous iron. It is widely distributed in peripheral organs, such as liver and kidney, and is also present in low amounts in the central nervous system. HAAO catalyzes the synthesis of quinolinic acid (QUIN) from 3-hydroxyanthranilic acid.

Nanjing BYabscience technology Co.,Ltd



QUIN is an excitotoxin whose toxicity is mediated by its ability to activate glutamate N-methyl-D-aspartate receptors. Increased cerebral levels of QUIN may participate in the pathogenesis of neurologic and inflammatory disorders. HAAO has been suggested to play a role in disorders associated with altered tissue levels of QUIN. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using 3HAO Monoclonal Antibody