



# AWAT2 Monoclonal Antibody

<b>Catalog No</b>	BYmab-05270
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	AWAT2 DC4 DGAT2L4 MFAT WS
<b>Protein Name</b>	Acyl-CoA wax alcohol acyltransferase 2 (EC 2.3.1.75) (Acyl-CoA retinol O-fatty-acyltransferase) (ARAT) (Retinol O-fatty-acyltransferase) (EC 2.3.1.76) (Diacylglycerol O-acyltransferase 2-like protein)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 260-340
<b>Specificity</b>	AWAT2 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	36kD
<b>Cell Pathway</b>	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
<b>Tissue Specificity</b>	Highly expressed in skin, where it is primarily restricted to undifferentiated peripheral sebocytes. Also expressed at lower level in other tissues except pancreas.
<b>Function</b>	catalytic activity:Acyl-CoA + a long-chain alcohol = CoA + a long-chain ester.,function:Acyltransferase that predominantly esterify long chain (wax) alcohols with acyl-CoA-derived fatty acids to produce wax esters. Wax esters are enriched in sebum, suggesting that it plays a central role in lipid metabolism in skin. Has no activity using decyl alcohol and significantly prefers the C16 and C18 alcohols. May also have 2-acylglycerol O-acyltransferase (MGAT) and acyl-CoA:retinol acyltransferase (ARAT) activities, to catalyze the synthesis of diacylglycerols and retinyl esters; however this activity is unclear in vivo.,similarity:Belongs to the diacylglycerol acyltransferase family.,tissue

**Nanjing BYabscience technology Co.,Ltd**



specificity: Abundant in tissues rich in sebaceous glands such as the preputial gland and eyelid. Highly expressed in skin, where it is primarily restricted to undifferentiated peripheral sebocytes. Also ex

## Background

This gene encodes an enzyme belonging to the diacylglycerol acyltransferase family. This enzyme produces wax esters by the esterification of long chain (or wax) alcohols with acyl-CoA-derived fatty acids. It functions in lipid metabolism in the skin, mostly in undifferentiated peripheral sebocytes. This enzyme may also have acyl-CoA:retinol acyltransferase activities, where it catalyzes the synthesis of diacylglycerols and retinyl esters. [provided by RefSeq, May 2010],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using AWAT2 Monoclonal Antibody