



ACO11 Monoclonal Antibody

Catalog No	BYmab-05276
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	ACOT11 BFIT KIAA0707 THEA
Protein Name	Acyl-coenzyme A thioesterase 11 (Acyl-CoA thioesterase 11) (EC 3.1.2.-) (Acyl-CoA thioester hydrolase 11) (Adipose-associated thioesterase) (Brown fat-inducible thioesterase) (BFIT)
Immunogen	Synthesized peptide derived from human protein . at AA range: 290-370
Specificity	ACO11 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	66kD
Cell Pathway	Mitochondrion matrix . Cytoplasm .
Tissue Specificity	Isoform 1 is predominantly expressed in skeletal muscle, liver, testis, stomach, spleen, lung and brain. Isoform 2 is predominantly expressed in kidney, uterus, hibernoma and white adipose tissue.
Function	function:Has acyl-CoA thioesterase activity towards medium (C12) and long-chain (C18) fatty acyl-CoA substrates.,induction:By cold exposure and repressed by heat exposure.,similarity:Contains 1 START domain.,similarity:Contains 2 acyl coenzyme A hydrolase domains.,tissue specificity:Isoform 1 is predominantly expressed in skeletal muscle, liver, testis, stomach, spleen, lung and brain. Isoform 2 is predominantly expressed in kidney, uterus, hibernoma and white adipose tissue.,
Background	This gene encodes a member of the acyl-CoA thioesterase family which catalyse the conversion of activated fatty acids to the corresponding non-esterified fatty

Nanjing BYabscience technology Co.,Ltd



acid and coenzyme A. Expression of a mouse homolog in brown adipose tissue is induced by low temperatures and repressed by warm temperatures. Higher levels of expression of the mouse homolog has been found in obesity-resistant mice compared with obesity-prone mice, suggesting a role of acyl-CoA thioesterase 11 in obesity. Alternative splicing results in transcript variants. [provided by RefSeq, Nov 2010],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658