



DDX42 Monoclonal Antibody

Catalog No	BYmab-05363
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DDX42
Protein Name	ATP-dependent RNA helicase DDX42 (EC 3.6.4.13) (DEAD box protein 42) (RNA helicase-like protein) (RHELP) (RNA helicase-related protein) (RNAHP) (SF3b DEAD box protein) (Splicing factor 3B-associated 1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 850-930
Specificity	DDX42 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	103kD
Cell Pathway	Cytoplasm . Nucleus .; [Isoform 2]: Nucleus, Cajal body . Nucleus speckle .
Tissue Specificity	Expressed in several cell lines (at protein level). Expressed in liver, lung, tonsil, thymus, muscle and pancreatic islets.
Function	function:ATP-dependent RNA helicase. Binds to partially double-stranded RNAs (dsRNAs) in order to unwind RNA secondary structures. Unwinding is promoted in the presence of single-strand binding proteins. Mediates also RNA duplex formation thereby displacing the single-strand RNA binding protein. ATP and ADP modulate its activity: ATP binding and hydrolysis by DDX42 triggers RNA strand separation, whereas the ADP-bound form of the protein triggers annealing of complementary RNA strands.,similarity:Belongs to the DEAD box helicase family. DDX42 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,subcellular location:Isoform 2 is present in Cajal bodies (CBs) and nuclear

Nanjing BYabscience technology Co.,Ltd



speckles.,subunit:Component of splicing factor SF3B which is composed of at least eight subunits; SF3B1/SAP155/SF3B155, SF3B2/SAP145/SF3B155, SF3B3/SAP130/SF3B130,

Background

This gene encodes a member of the Asp-Glu-Ala-Asp (DEAD) box protein family. Members of this protein family are putative RNA helicases, and are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. Two transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658